Spinal and limb abnormalities in adolescents with intellectual disabilities

Jin-Ding Lin^{a,*}, Pei-Ying Lin^a, Lan-Ping Lin^b, Chia-Im Lai^c, Yii-Rong Leu^d, Chia-Feng Yen^e, Shang-Wei Hsu^f, Chi-Ming Chu^a, Chia-Ling Wu^g, Cordia M. Chu^h

^aSchool of Public Health, National Defense Medical Center, Taipei, Taiwan

^bGraduate Institute of Life Sciences, National Defense Medical Center, Taipei,

Taiwan

^cBranch of Medical Affair Management, Tri-Service General Hospital, Taipei,

Taiwan

^dInstitute of Gerontology, Heidelberg University, Germany

^eDepartment of Social Welfare, National Chung Cheng University, Chiayi, Taiwan

^fGraduate Institute of Healthcare Administration, Asia University, Taichung, Taiwan

^gResearch Center for Intellectual Disabilities Taiwan, Chung-Hua Foundation for

Persons with Intellectual Disabilities, Taipei County, Taiwan

^hCenter for Environment and Population Health, Griffith University, Brisbane,

Australia

*Corresponding author: Jin-Ding Lin, School of Public Health, National Defense

Medical Center, Taipei, Taiwan

No. 161, Min-Chun East Road, Section 6, Nei-Hu, Taipei, Taiwan

Tel: 886-2-87923100#18447

Fax: 886-2-87923147

E-mail: a530706@ndmctsgh.edu.tw

Abstract

There are not many studies pertaining to the spinal or limb abnormalities in people with intellectual disabilities, without a clear profile of these deformities of them, efforts to understand its characters and improve their quality of life will be impossible. Therefore, this paper aims to describe the prevalence and related factors of spinal and limb abnormalities in adolescents with intellectual disabilities. The subjects who participated in health examinations as they enrolled into special schools at the first year, a total of 822 aged 15-18 years adolescents with ID were recruited to this study. The results showed that there were 14.5% and 8.5% cases had spinal and limb abnormalities based on the physician's observation and X-ray test. Factors of BMI level and limb abnormalities were significantly predicted the spinal abnormality occurrence in those adolescents with ID. Gender, disability level and have a spinal abnormality were variables that can statistically correlate to limb abnormality condition. The study highlights that in order to ensure people with intellectual disabilities receive an appropriate quality of care, it is important to have a precise understanding of the ways in which the needs of them who have spinal or limb deformities differ from the sole intellectual disability and the general population as a whole.

Keywords: Intellectual disability, spinal abnormality, limb abnormality, health exam

Introduction

There are not many literature pertaining to the spinal or limb abnormalities in people with intellectual disability (ID) in the previous decades. One of many earlier studies, G. B. Solitare described "The spinal cord of the Mongol" in 1969, he reported four autopsies on Mongols (now Down syndrome) to examine their spinal cord lesion (Solitare, 1969). Smith, Schindeler, Elbualy and Shear (1970) indicated that limb abnormalities are occasionally found in populations of ID individuals. Recent studies, Tangerus, Hestnes, Sand and Sunndalsfoll (1990) indicated that there is a significant increase in degenerative changes in the upper part of the cervical spine in Down's syndrome. Maclachlan et al. (1993) also concerned that adults with Down syndrome are at high risk of high prevalence of degenerative disease of the cervical spine and consideration should be given to this diagnosis in the appropriate clinical setting.

The consequences of spinal or limb abnormalities in people with ID are complicated. Mori et al. (2005) examined the correlation between spinal characteristics and respiratory function in patients with severe physical disabilities and ID. These observations suggest that the spinal abnormalities affect respiratory patterns in a restrictive manner and increase the respiratory change rate during tidal breathing. Lutkenhoff & Oppenheimer (1997) described the adolescents with spina bifida "Young people with spina bifida with have many questions about themselves and their futures which they may not want to ask of their parents or doctors. Even if they do ask, there is likelihood that the person asked won't know the answer". However, without a clear profile of spinal or limb deformities in people with ID, efforts to understand its characters and improve their quality of life will be impossible. Therefore, the purposes of this paper were to describe the prevalence and related factors of spinal and limb abnormalities in adolescents with ID.

Methods

This sample group is part of a retrospective, cross-sectional study that examines the health examination results for children and adolescents with ID (Lin et al., 2009; Lin et al., 2010a, 2010b). Subjects were recruited from three special educational schools, the adolescents with ID who participated in health examinations as they enrolled into schools at the first year. The examination includes body physical exam, biochemical (blood, urine and stool specimen) and X-ray check-up. The study was designed to analyze all the health examination charts which included all eligible aged 15-18 years adolescents with ID from all three special schools. Research ethical approval and written informed consent were obtained from all the study special schools.

The present study aims to analyze the conditions of spinal and limb abnormalities and to examine the determinants of these two characteristics. There were 822 participants data were analyzed, those data included demographic characteristics (age and gender), disability condition (type and level), body mass index (BMI; kg/m²), hands, feet and spinal abnormalities. Hands and feet abnormalities (called limb abnormalities) were based on the medical physicians' observations, and X-ray was used to determine the spinal abnormalities. Finally, those abnormalities which showed in medical charts were based on the diagnosis results of medical physicians in the study. Data were analyzed by SPSS 14.0. Analyses included frequency distributions and percentage for the demographic characteristics and the prevalence of spinal and limb abnormalities among adolescents with ID. Chi-square tests were obtained for the relationship between demographic characteristics, BMI value and spinal and limb abnormalities. Logistic regression procedures were used to examine the risks of having spinal and limb abnormalities.

Results

A total of 822 aged 15-18 years adolescents with ID were recruited to this study. The mean age of study subjects was 15.69±0.75 years old, 60.7% subjects were boys and 39.3% were girls. In our samples, most of the people with ID had a moderate level of disability (52.7%) and severe and profound disability accounting for 29.6%. There was nearly seventy percent of the study subjects were ID solely while 30.5% were affected by multiple disabilities (ID accompanied with other disabilities) as well. With regard to the physical figure of the subjects, the results of BMI analysis indicated their average score was 22.2 and 34.1% subjects were normal, 13.7% were overweight, 24.9% were obese and 27.3% were underweight. The physical observation and X-ray test results showed that there were 14.5% and 8.5% cases had spinal and limb abnormalities in the study.

Table 2 and table 3 analyzed chi-square relation between personal characteristics and spinal or limb abnormalities in the bivariant analysis. The results showed BMI was significantly correlated to spinal abnormalities (p<0.001). Underweight adolescents with ID (prevalence=22.9%) were more likely to have spinal abnormalities than the other BMI level groups. Those factors of gender, disability type and level were not correlated to spinal abnormalities in adolescents with ID. However, the factors of disability type and disability level were statistically correlated to limb abnormalities in adolescents with ID. Those adolescents with multiple disabilities (abnormal prevalence=19.3%) were more than 5 times to have limb abnormalities than the ID solely cases (abnormal prevalence=3.7%). In

addition, the more serious of the disability level, the more prevalent of the limb abnormalities in adolescents with ID in the study (p<0.001). Table 4 also found that spinal abnormality was significantly correlated to limb abnormality in the study subjects (p<0.001).

Table 5 tested the logistic regression model of spinal abnormality occurrence of adolescents with ID. The results showed the BMI level (p=0.035) and limb abnormalities (p<0.001) were significantly predicted the spinal abnormality occurrence in those adolescents with ID (p<0.05). The spinal abnormality cases tends to be more prevalent in the underweight group than in the normal weight group (OR=1.82, 95%CI=1.04-3.18), and the limb abnormality cases were more likely to have spinal abnormalities than those adolescents do not have limb abnormalities (OR=6.12, 95%CI=3.15-11.9). With respect to the risk factors of limb abnormality occurrence in adolescents with ID, we found that the factors of gender, disability level and spinal abnormality were variables that can significantly predict they have a limb abnormality condition after controlling factors of disability type, age and BMI in the study (table 6). Those boys adolescents were less chance to have a limb abnormality than girl cases (OR=0.47, 95% CI=0.25-0.90). However, those subjects with multiple disabilities (OR=3.63, 95% CI=1.72-7.58) and have a spinal abnormality (OR=5.90, 95% CI=3.01-11.56) were more likely to have a limb abnormality in the study.

Discussion

Traditionally, the medical care needs and problems of persons with ID living in the general community have received limited attention in healthcare systems (Minihan & Dean, 1990; van Schrojenstein Lantman-de Valk et al., 1997; Lin et al., 2007; Lin, 2009; Lin, Yen, Wu, & Kang, 2009). Many studies have demonstrated that individuals with ID experience poorer health (Li, Wu, Lee, 2003; Lin, Yen, Li, Wu, 2005; Yen, Loh, & Lin, 2009; Lin et al., 2009; Lin et al., 2010a, 2010; Yen & Lin, 2010) and use more medical care services than the general population (Lin, Loh, Yen, Li, & Wu, 2006; Hsu et al., 2009; Lin, Lin, Yen, Loh, & Chwo, 2009; Yen, Lin, Loh, Shi, & Shu, 2009). Most of them experience barriers in accessing health services and supports for their health needs (Lin et al., 2005; Lin, Hsu, Yen, Chou et al., 2009). To ensure people with ID can enjoy the same health right as the general population, the health care system should take further steps to develop an appropriate monitoring system and health services for them (Lin et al., 2005). This paper aims to examine the prevalence and related factors of spinal and limb abnormalities in adolescents with ID. We found that there were 14.5% and 8.5% cases had spinal and limb abnormalities in the study based on physician's observation and X-ray test. Factors of BMI level and limb

abnormalities were significantly predicted the spinal abnormality occurrence in those adolescents with ID. Gender, disability level and have a spinal abnormality were variables that can statistically correlated to limb abnormality condition.

Smith, Schindeler, Elbualy and Shear (1970) stated that sometimes the limb abnormalities may be so characteristic of a syndrome that the diagnosis can be made directly. In most instances, the limb abnormality is not specific and it is only diagnostic when the total clinical complex is examined. They highlighted the limb abnormalities in people with ID may frequently suggest a specific diagnosis or syndrome. The present study highlights the high abnormal rate in spinal and limb among adolescents with ID needed to pay much attention in healthcare system. Many studies have provided the opportunities to therapy or correct of spinal or limb abnormalities. Burch, Clegg and Bailey (1987) demonstrated that electronic switches can be used effectively as therapeutic tools to correct improper posture of the individuals during functional daily activities. Behavior therapy or physical therapies are also good tools to promote independent ambulation in people with ID (Horton & Taylor, 1989). Generally, physical rehabilitation following spinal cord injury-related paralysis or abnormalities has traditionally focused on teaching compensatory techniques, thus enabling the individual to achieve daily function despite significant neurological deficits (Sadowsky & McDonald, 2009). They supported the idea of utilizing physical activity and exercise are not luxury, but needed therapeutic intervention to manage the condition such as spinal cord injury, and eventually other disorders of the CNS like stroke and CP.

Two limitations in the present study need to be pointed out. First, we only recruited cases who studying in special schools; it may not represent the full situations of adolescents with ID. Second, the health exam results lack double check of another health setting will possible affect the results. However, this study provides a general profile of physical deformities among adolescents with ID. To ensure that people with ID receive an appropriate quality of care, it is important to have a precise understanding of the ways in which the needs of them who have spinal or limb deformities differ from the sole ID and the general population as a whole.

Acknowledgements

This research was financially supported by Bureau of Health Promotion, Department of Health, Executive Yuan, Republic of China (Taiwan) (DOH97-HP-1601). We would also like to thank the special educational school's staffs who support this study.

References

- Burch, M. R., Clegg, J. C., & Bailey, J. S. (1987). Automated Contingent reinforcement of correct posture. *Research in Developmental Disabilities*, 8, 15-20.
- Horton, S. V., & Taylor, D. C. (1989). The use of behavior therapy and physical therapy to promote independent ambulation in a preschooler with mental retardation and cerebral palsy. *Research in Developmental Disabilities*, 10, 363-375.
- Hsu, S. W., Lin, Y. W., Chwo, M. J., Huang, H. C., Yen, C. F., Lin, L. P., Wu, J. L., & Lin, J.D. (2009). Emergency department utilization and determinants of use by 0 to 6-year-old children with disabilities in Taipei. *Research in Developmental Disabilities*, 30, 774-781.
- Lin, J. D. (2009). Population with intellectual disability based on 2000-2007 national registers in Taiwan: Age and gender. *Research in Developmental Disabilities*, 30, 294-300.
- Lin, J. D., Hsu, S. W., Yen, C. F., Chou, Y. T., Wu, J. L., Chu, C., & Loh, C. H. (2009). Role of the general practitioners in the provision of health care services for people with intellectual disabilities: A national census in Taiwan. *Journal of Applied Research in Intellectual Disabilities*, 22, 582-591.
- Lin, J. D., Lin, P. Y., Chen, L. M., Fang, W. H., Lin, L. P., Loh, C. H. (2010a). Serum glutamic oxaloacetic transaminase (GOT) and glutamic pyruvic transaminase (GPT) levels in children and adolescents with intellectual disabilities. *Research in Developmental Disabilities*, 31, 172-177.
- Lin, J. D., Lin, P. Y., Lin, L. P., Hsu, S. W., Loh, C. H., Yen, C. F., Fang, W. H., Chien, W. C., Tang, C. C., & Wu, C. L. (2010b). Prevalence and associated risk factors of anemia in children and adolescents with intellectual disabilities. *Research in Developmental Disabilities*, 31, 25-32.
- Lin, J. D., Lin, P. Y., Lin, L. P., Hsu, S. W., Yen, C. F., Fang, W. H., Wu, S.R., Chien, W. C., Loh, C. H., & Chu, C. (2009). Serum uric acid, hyperuricemia and body mass index in children and adolescents with intellectual disabilities. *Research in Developmental Disabilities*, 30, 1481-1489.
- Lin, J. D., Lin, Y. W., Yen, C. F., Loh, C. H., & Chwo, M. J. (2009). Received, understanding and satisfaction of National Health Insurance premium subsidy scheme by families of children with disabilities: A census study in Taipei City. *Research in Developmental Disabilities*, 30, 275-283.
- Lin, J. D., Loh, C. H., Yen, C. F., Li, C. W., Chwo, M. J., & Wu, J. L. (2007). Medical care services for people with intellectual disabilities living in the general community: A cross-sectional survey of inpatient care utilization in Taiwan, 2001.

- Disability and Rehabilitation, 29, 1411-1416.
- Lin, J. D., Loh, C. H., Yen, C. F., Li, C. W., & Wu, J. L. (2006). Rehabilitation service utilization and determinants among people with an intellectual disability: Preliminary findings in Taiwan. *Disability & Rehabilitation*, 28, 1499-1506.
- Lin, J. D., Wu, J. L., & Lee, P. N. (2003). Health care needs of persons with intellectual disabilities in institutions in Taiwan: Outpatient utilization and implications. *Journal of Intellectual Disability Research*, 47, 169-180.
- Lin, J. D., Yen, C. F., Chwo, M. J., Tung, H. J., & Lee, T. N. (2005) Primary health care for people with an intellectual disability: A mission impossible? *Journal of Medical Sciences*, 25, 109-118.
- Lin, J. D., Yen, C. F., Li, C. W., & Wu, J. L. (2005). Health, healthcare utilization and psychiatric disorder in people with intellectual disability in Taiwan. *Journal of Intellectual Disability Research*, 49, 86-94.
- Lin, J. D., Yen, C. F., Wu, J. L., & Kang, S. W. (2009). National disability registers report on causes of intellectual disability in Taiwan: 2000-2007. *Research in Developmental Disabilities*, 30, 301-307.
- Lutkenhoff, M., & Oppenheimer, S. G. (1997). Spinabilities: A young person's guide. Bethesda: Woodbine House.
- Maclachlan, R. A., Fidler, K. E., Yeh, H., Hodgetts, P. G., Pharand, G., & Chaum, M. (1993). Cervical spine abnormalities in institutionalized adults with Down's syndrome. *Journal of Intellectual Disability Research*, 37, 277-285.
- Minihan, P. M., & Dean, D. D. (1990). Meeting the needs of health services for persons with mental retardation living in the community. *American Journal of Public Health*, 80, 1043-1048.
- Mori, N., Kurosawa, H., Hiramoto, A., Matumoto, K., Sano, Y., Fukuda, E., Fujii, H., & Kohzuki, M. (2005). Relationships between respiratory function and spinal characteristics in patients with severe motor and intellectual disabilities syndrome. *Chest*, 128, 355S.
- Sadowsky, C. L., & McDonald, J. W. (2009). Activity-based restorative therapies concepts and applications in spinal cord injury-related neurorehabilitation. *Developmental Disabilities Research Reviews*, 15, 112-116.
- Smith, G. F., Schindeler, J., Elbualy, S., & Shear, C. (1970). Limb abnormalities and mental retardation. *Journal of Mental Deficiency Research*, 14, 319-341.
- Solitare, G. B. (1969). The spinal cord of the Mongol. *Journal of Mental Deficiency Research*, 13, 1-7.

- Tangerud, A., Hestnes, A., Sand, T., Sunndalfoll, S. (1990). Degenerative changes in the cervical spine in Down's syndrome. *Journal of Intellectual Disability Research*, 34, 179-185.
- van Schrojenstein Lantman-de Valk, H. M., Metsemakers, J. F., Soomers-Turlings, M. J., Haveman, M. J., & Crebolder, H. F. (1997). People with intellectual disability in general practice: case definition and case finding. *Journal of Intellectual Disability Research*, 41, 373-379.
- Yen, C. F., & Lin, J. D. (2010). Factors for healthy food or less-healthy food intake among Taiwanese adolescents with intellectual disabilities. *Research in Developmental Disabilities*, 31, 203-211.
- Yen ,C. F., Lin, J. D., Loh, C. H., Shi, L., & Shu, S. W. (2009) Determinants of Prescription drug use by adolescents with intellectual disabilities in Taiwan. *Research in Developmental Disabilities*, 30, 1354-1366.
- Yen, C. F., Loh, C. H., & Lin, J. D. (2009). Prevention of mental health problems in people with intellectual disabilities. *Current Opinion in Psychiatry*, 22, 447-451.

Table 1. Subject's demographic characteristics and their body deformities

Characteristics	n	%	Mean±S.D. (Range)
Condor (N-922)			
Gender (N=822)	400	<i>c</i> 0.7	
Boys	499	60.7	
Girls	323	39.3	
Age (years) (N=822)			15.69±0.75 (15-18)
Disability type (N=822)			
ID	571	69.5	
Multiple [*]	251	30.5	
Disability level (N=822)	231	30.3	
Mild	80	9.7	
Moderate	433	52.7	
Serve			
	243	29.6	
Profound	66	8.0	
$BMI(kg/m^2)$ (N=812)			22.19±5.55 (8.7-50.4)
Underweight	222	27.3	
Normal	277	34.1	
Overweight	111	13.7	
Obesity	202	24.9	
Spine (N=812)			
Normal	694	85.5	
Abnormal	118	14.5	
Limb (N=701)			
Normal	641	91.4	
Abnormal	60	8.6	

^{*}ID accompanied with other disabilities

Table 2. Relation of subject's characteristics and spine abnormalities

Variable	Spine		χ^2	p value
	Normal	Abnormal		
	(percent)	(percent)		
Gender (N=812)			0.11	0.737
Boys	418 (60.2)	73 (14.9)		
Girls	276 (86.0)	45 (14.0)		
Disability type (N=812)			0.38	0.537
ID	490 (86.0)	80 (14.0)		
Multiple	204 (84.3)	38 (15.7)		
Disability level (N=812)			2.46	0.482
Mild	71 (88.8)	9 (11.3)		
Moderate	370 (85.5)	63 (14.5)		
Severe	202 (86.0)	33 (14.0)		
Profound	51 (79.7)	13 (20.3)		
BMI (N=806)			21.45	< 0.001
Underweight	168 (77.1)	50 (22.9)		
Normal	244 (88.7)	31 (11.3)		
Overweight	92 (82.9)	19 (17.1)		
Obese	185 (91.6)	17 (8.4)		

Table 3. Relation of subject's characteristics and limb abnormalities

Variable	Limb		χ^2	p value
	Normal	Abnormal		
	(percent)	(percent)		
Gender (N=701)			2.45	0.118
Boys	387 (92.8)	30 (7.2)		
Girls	254 (89.4)	30 (10.6)		
Disability type (N=701)			46.34	< 0.001
ID	465 (96.3)	18 (3.7)		
Multiple	176 (80.7)	42 (19.3)		
Disability level (N=701)			43.16	< 0.001
Mild	67 (98.5)	1 (1.5)		
Moderate	355 (95.9)	15 (4.1)		
Severe	178 (85.6)	30 (14.4)		
Profound	41 (74.5)	14 (25.5)		
BMI (N=691)			7.19	0.066
Underweight	173 (89.2)	21 (10.8)		
Normal	216 (93.9)	14 (6.1)		
Overweight	87 (90.6)	9 (9.4)		
Obese	164 (95.9)	7 (4.1)		

Table 4. Relation of spinal and limb abnormalities (N=691)

X7 ' 1 1	S :	. , ,	\mathbf{v}^2	p value
Variable	Spine	Spine		
	Normal (percent)	Abnormal (percent)		
Limb			32.89	< 0.001
Normal	555 (87.4)	80 (12.6)		
Abnormal	33 (58.9)	23 (41.1)		

Table 5. Logistic regression model of spinal abnormalities (N=685)

Variable	β	S.E.	OR	95% CI	p value
Constant	-2.38	2.33	0.09		0.307
Gender					
Girls			1.00		
Boys	0.16	0.23	1.17	0.74-1.85	0.500
Age	-0.01	0.15	1.00	0.74-1.33	0.972
Disability type					
ID			1.00		
Multiple	-0.48	0.31	0.62	0.34-1.14	0.122
Disability level					
Mild			1.00		
Moderate	0.38	0.44	1.47	0.63-3.44	0.377
Severe	0.18	0.48	1.20	0.47-3.10	0.706
Profound	0.88	0.59	2.40	0.75-7.64	0.139
BMI					
Normal			1.00		
Underweight	0.60	0.28	1.82	1.04-3.18	0.035
Overweight	0.44	0.35	1.55	0.79-3.05	0.206
Obese	-0.26	0.34	0.77	0.40-1.51	0.451
Limb					
Normal			1.00		
Abnormal	1.81	0.34	6.12	3.15-11.90	< 0.001

Table 6. Logistic regression model of limb abnormalities (N=685)

Variable	β	S.E.	OR	95% CI	p value
Constant	-7.43	3.25	0.001		0.022
Gender					
Female			1.00		
Male	-0.75	0.33	0.47	0.25-0.90	0.022
Age	0.20	0.20	1.23	0.83-1.81	0.306
Disability type					
ID			1.00		
Multiple	1.29	0.38	3.63	1.72-7.58	0.001
Disability level					
Mild			1.00		
Moderate	0.62	1.06	1.86	0.23-14.90	0.557
Severe	1.42	1.08	4.13	0.50-33.99	0.188
Profound	1.50	1.14	4.48	0.48-41.72	0.188
BMI					
Normal			1.00		
Underweight	0.28	0.41	1.32	0.60-2.93	0.493
Overweight	0.28	0.49	1.32	0.50-3.45	0.575
Obese	-0.41	0.50	0.67	0.25-1.77	0.415
Spine					
Normal			1.00		
Abnormal	1.78	0.34	5.90	3.01-11.56	< 0.001