# SIMPLE PREDATOR-PREY INTERACTIONS CONTROL DYNAMICS

# IN A PLANKTON FOOD WEB MODEL

- 5 Roger Cropp<sup>1</sup> and John Norbury<sup>2</sup>
- 6 <sup>1</sup> Centre for Environmental Systems Research, Griffith School of Environment, Griffith
- 7 University, Nathan, Queensland, 4111, Australia.
- 8 <sup>2</sup> Mathematical Institute, University of Oxford, 24-29 St Giles, Oxford, OX1 3LB, UK.

# **ABSTRACT**

A plankton food web model is analysed using interaction parameter values appropriate to the upper mixed layer of the high latitude oceans. The dynamics of this four-variable system are analysed in terms of the dynamics of much simpler two-variable predator-prey subsystems. Thus, the food web's robust, periodic, four-dimensional dynamics are explained by means of two-dimensional spirals and limit cycles. These dynamical subsystems are coupled by means of an omnivore that transfers control of the dynamics between the two predator-prey subsystems. The food web may substantially decouple the predator-prey subsystems so that the oscillating phytoplankton/zooplankton blooms exhibit population collapses when bacterial 'breathers' briefly dominate after growing dramatically from low background levels. This regular bloom/breather behaviour becomes benignly chaotic when the system is mildly forced by the annual cycle of the sun's irradiance.

1 Key words: Plankton modelling
2 Population dynamics
3 Kolmogorov systems
4
5

## 1. INTRODUCTION

Marine plankton ecosystems are being increasingly recognised as a potentially important influence on global climate (Le Quere et al. 2005). Marine plankton ecosystems are an integral component of biogeochemical cycling in the oceans and may have important climate effects due to their ability to draw down carbon dioxide from the atmosphere and store it in the deep ocean (Falkowski et al. 2000) and for their potential to affect cloud formation by producing gases that are eventually transferred to the atmosphere (Charlson et al. 1987). The evaluation of the potential of both of these climatically important processes to affect climate depends critically on understanding the interactions between members of ocean plankton ecosystems, that is, the dynamics of the ecosystem.

Generally, the veracity of plankton ecosystem models is evaluated by numerically integrating the models and determining how well they can reproduce observed variables (Le Quere et al. 2005). Typically, ecosystem models are 'calibrated' (the values of parameters describing feeding rates, etc are selected from within a measured range) by comparing model outputs with measured data until the best 'fit' is obtained. The models are then 'validated' by comparing the predictions of the calibrated model with an independent data set. However, a survey of plankton ecosystem models by Arhonditsis and Brett (2004) noted that 95% of plankton ecosystem modelling publications did not report on the model's performance in reproducing all state variables and only 30% quantified the comparison between observed and modelled data. Arhonditsis and Brett observed that about half of modelling papers report that the models were subjected to sensitivity analysis, with a similar proportion validated with independent data.

1 Complex ecosystem models present special problems for this approach, as they often have

more parameters than can be constrained by data (Matear 1995).

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

2

The application of plankton ecosystem models in climate modelling has placed additional demands on modellers to develop ever more complex models. (Here we use 'complex' to explicitly mean 'large and complicated', rather than in the context of 'complex systems science' where it is often interpreted to mean irreducible.) Ecosystem models are commonly expressed mathematically as coupled ordinary differential equations (odes) and are often written in a single 'currency', for example in terms of a limiting nutrient such as atomic nitrogen. However, the demands of climate modelling for resolution of the fate of climaticallyimportant gases such as carbon dioxide and dimethyl sulphide in seawater means that these models must also include currencies of carbon and sulphur (Cropp et al. 2004). Similarly, there is increasing evidence that the utilisation of available nitrogen may be limited by the availability of iron, suggesting that iron may also need to be included. If bacterial processes are important, then more than one form of nitrogen may need to be included. The resolution of important processes such as the sinking and export to the deep ocean may require plankton functional types to be included (and consequently even more nutrients explicitly represented), as biogeochemical cycling in marine systems appears closely coupled to particular plankton groups (Anderson 2005). Complex plankton ecosystem models may therefore include 70 state variables (Baretta et al. 1995, Lancelot et al. 2000, Arhonditsis & Brett 2004).

21

22

23

24

25

Mathematical analysis of the dynamics of complex ecosystem models becomes especially difficult when there are three or more state variables and numerical methods must often be employed. Sophisticated numerical integrators are now available that can approximate the dynamics and long-term states of complex ode models; however, vagaries in the

implementation of numerical solvers can lead to different results for the same model implementation (Seppelt & Richter 2005), emphasising the importance of more qualitative assessments of ecosystem models. Sensitivity and uncertainty analysis can provide some increased level of confidence in model results by evaluating the sensitivity of the results to variations in the input parameter values. However, sensitivity analysis techniques require many model evaluations, and the computational expense of a global sensitivity analysis of a complex

ecosystem model can be prohibitive.

There is a rich literature attesting to the sensitivity of the dynamics of ode models of ecosystems to small variations in parameter values when, for example, steady states may become oscillatory (Edwards & Brindley 1999). This sensitivity to small parameter variations is especially important for ecosystem models that are coupled to global climate models (Earth System Models) to incorporate ocean biogeochemistry into global climate predictions. The warming of the oceans associated with climate change can affect the parameters of plankton ecosystems in many ways. When simulating ecosystem responses to climate change then, the fact that the models can reproduce current states is no guarantee that the models will correctly predict future climates – the susceptibility of the dynamics of model ecosystems to parameter variations is well-known, and minor parameter changes can lead to significant changes in dynamical behaviours (Kuznetsov & Rinaldi 1996).

An analytic explication of the factors driving the ecosystem model dynamics can therefore provide a valuable adjunct to numerical techniques. However, nonlinear dynamical systems theory is mostly established for ode models with only two state variables; few theorems extend to three or more variables. An early but often overlooked example of the application of nonlinear dynamical systems theory to ecosystem models is the work of Kolmogorov (1936) in

which he developed conditions that ensure that a particular class of two-dimensional predator-

prey models, that have come to be called "Kolmogorov systems", have either a stable

equilibrium or a stable limit cycle. May (1973) discussed the ecological interpretation of

Kolmogorov's theorem, that oscillations in ecosystems either constantly repeat periodically or

die out to a stable equilibrium, and noted that it applied to many ecological models then in use.

Kolmogorov systems have attracted substantial mathematical interest as they include many

types of predator-prey models (Huang & Zhu 2005), but they have been largely ignored by

ecologists (Holling 1973).

An important issue in ecology is to understand the dynamics of complex trophic systems as a result of the direct interactions between pairs of species. Predator-prey interactions (or, more generally, the interactions between consumers and their resources) are the defining modules in aquatic food webs but it is an open question whether analyses of small subsets of species can provide insights that are relevant to larger communities and ecosystems. In this paper, we explore the insights into the factors controlling the dynamics of complex ecosystems that may be gained by breaking a complex ecosystem model down into its constituent smaller food-web models. Neutel et al. (2002, 2007) used an analogous approach to categorise the properties of real soil micro-organism food-webs, but their subsystems were not self-sustaining. In contrast, we consider only subsystems that are self-sustaining, that is each subsystem we consider could survive as an autonomous unit in the absence of the other organisms in the full food web.

We examine a moderate complexity marine plankton ecosystem model based on one developed

by Moloney et al. (1986) that we have calibrated with parameter values demonstrated to be

valid for simulating plankton dynamics in the Barents Sea (Gabric et al. 1999). This model is

based on commonly used functional relationships between bacteria (B), zooflagellates (F),

- phytoplankton (P) and zooplankton (Z), and includes a limiting nutrient (N). The key features
- 2 of this model are that it is a 'Kolmogorov system' and that it conserves mass.

- 4 Many contemporary ecosystem models (Spitz et al. 2001, Franks 2002, Vallina et al. 2008)
- 5 may be classed as Kolmogorov systems, as they are of the general form:

$$\dot{u}_i = f_i (u_1, u_2, K, u_n) u_i, \quad i = 1, 2, K, n,$$
 (1)

7 where  $\dot{u} = \frac{du}{dt}$  for t > 0, and the functions  $f_i$  are bounded and continuously differentiable in

8 their variables  $u_1, u_2, K, u_n$ . We make the distinction here between a Kolomogorov system,

9 which is any system that may be written in the form of (1), and Kolmogorov *conditions*, which

describe the conditions on a Kolmogorov system that ensure ecologically realistic dynamics

(Kolmogorov 1936). We observe, as did May (1973), that Kolmogorov's conditions may be

12 relaxed whilst still ensuring reasonable dynamics.

13

15

16

17

18

19

20

21

22

23

10

11

14 The  $f_i$  in equation (1) describe the net growth and mortality of each species, functional type,

guild or trophic level, that is  $f_i = (growth - predation - mortality)_i$ , and are often nonlinear

functions of  $u_1, u_2, K, u_n$ . There are many options for the  $f_i$  but in accordance with

Kolmogorov's (1936) criteria for ecologically realistic dynamics, we choose simple

ecologically realistic functions such that the autotrophs are the only variables that can grow at

low population concentrations (i.e.  $f_a(0,0) > 0$  and  $f_p(0,0) < 0$ ) where a denotes autotrophs

and p denotes predators and all other non-autotrophs. Further, increasing population density

while maintaining constant ratios of the predators and prey has different effects for predators

and prey. The prey is less able to thrive while the predator is more able to thrive as populations

increase. In mathematical parlance this is described as  $u \cdot \nabla f_a < 0 < u \cdot \nabla f_p$  (May 1973).

1 Ecological realism may also be seen with relation to the  $f_a = 0$  and  $f_p = 0$  isoclines (for

example, Figure 1). Here, we want the  $f_p$  isocline to intersect the  $f_a$  isocline only once, and to

3 intersect in the manner of the diagram, where for positive  $\alpha$ ,  $\beta$ ,  $\gamma$  we have that  $f_a(\alpha, 0) = 0$ ,

4  $f_a(0,\beta) = 0$  and  $f_p(\gamma,0) = 0$  imply  $\alpha > \gamma$  as per Rescigno and Richardson (1967) and May

(1973). Note the different scales of intersections in the first two diagrams of the upper panel

compared to that of the third: this reflects the different scale of the B processes.

7

11

12

13

14

15

16

17

18

19

20

21

5

6

8 Conservation of mass, or 'closure', depends on there being a currency in terms of which we 9 can measure the concentrations of both the predator and the prey. Here we call this nutrient, 10 and closure means that the total amount of nutrient in the system (the sum of both inorganic

and organic forms) remains constant for all time in spite of the complicated interactions

between biota. Such conservation of mass is explicit in many models of plankton dynamics

(Franks 2002, Gibson et al. 2005), and we also observe that some models that do not explicitly

conserve mass fit observed data best when the nutrient uptake and loss fluxes approximately

balance (Spitz et al. 2001). Conservation of mass within the mixed layer is also a commonly

observed property of marine planktonic ecosystems, that typically cycle nutrient very tightly

within the mixed layer of the ocean, and is referred to as regenerated production (Dugdale &

Goering 1967). Pragmatically, enforcing conservation of mass in an ecosystem model that is

not a Kolmogorov system can in many cases allow it to be written as a Kolmogorov system,

allowing the extension of this approach to many other models for the special case where

nutrient inputs equal nutrient losses.

22

23

24

Mathematically, conservation of mass means that, when the scaled variables  $u_1, u_2, K$ ,  $u_n$  are

expressed in terms of the nutrient (N) that the variables either feed on or decay into, the total

1 amount of nutrient is conserved and does not change with time; that is

2  $u_1 + u_2 + K + u_n + N = N_T$ . This provides a further equation to our Kolmogorov system;

3 although this final equation is not in Kolmogorov form, it is technically redundant as it may be

4 derived directly from the closure of mass condition:

$$N = N_T - u_1 - u_2 - \mathbf{K} - u_n \iff N = -i\mathbf{k} - i\mathbf{k} - \mathbf{k} - i\mathbf{k}$$
 (2)

6 For a suitably scaled model where the total mass is one unit, the amount of inorganic nutrient

7 (N) present at any time is given by  $N = 1 - u_1 - u_2 - K - u_n$ ; when  $\dot{N} > 0$  on N = 0 for  $u_p > 0$ 

this condition defines an ecologically feasible 'state space' where  $0 \le u_1, u_2, K$ ,  $u_n \le 1$  and

 $u_1 + u_2 + K + u_n \le 1 \Leftrightarrow N \ge 0$ . The dynamics of the system are then confined to the part of a

multi-dimensional Cartesian co-ordinate space where each axis represents the (suitably scaled)

concentration of the predator or prey and all the variables are positive and less than one. The

condition  $\sum_{i=1}^{n} u_i \le 1$  provides a tighter 'lid' on the dynamics. As described by May (1973), such

13 Kolmogorov systems are realistic descriptions of basic ecological models in that populations

oscillate for ever or their oscillations gradually decay to come to a stable equilibrium.

15

16

17

18

19

20

21

22

14

8

9

10

11

12

We shall consider a complex food-web system that involves the interconnection of two

archetypal predator-prey subsystems, each of which is a Kolmogorov system. The predator-

prey subsystems are linked by the omnivorous properties of one of the autotrophs. We observe

that our moderate-complexity food web has a very robust stable limit cycle with successive

blooms of the biota. The limit cycle is comprised of decaying oscillations ('blooms') of one

predator-prey interaction interspersed with occasional outbreaks, that we call 'breathers', of

bacteria that are rapidly controlled by their predator. The most interesting aspect of our model,

23 however, is that it demonstrates that the dynamical behaviours of certain realistic complex

1 models can be inferred from the dynamical behaviours of the predator-prey subsystems that

2 comprise them.

#### 2. THE PLANKTON MODEL

numbers of individuals or biomass.

We consider, as an explicit example, a model based on the plankton ecosystem model developed by Moloney et al. (1986). This four-dimensional closed Kolmogorov system was developed from consideration of the allometry of plankton, and has been applied in studies to simulate the response of marine ecosystems to climate change and to model their potential to mitigate the extent of global warming (Gabric et al. 2003). The version of the model we consider has four trophic groups; bacteria (B), zooflagellates (F), phytoplankton (P) and zooplankton (Z) and is written in a currency of the limiting nutrient nitrogen (N). The model, which we shall refer to as the BFNPZ model, is comprised of four equations that define the concentration of nutrient contained in each of the biota at any time. The populations of the biota are therefore described in terms of the atomic nitrogen that they contain, rather than

Each of the model equations is composed of terms that represent each organisms' growth, from consuming inorganic nutrient or other organisms; losses to grazing by their predators; and mortality, including both senescence and losses to predation by higher predators that are only implicitly represented in the model (i.e. are not represented by an equation explicitly describing how their population varies over time). These processes are represented in the model using functional forms commonly utilised in ecosystem models: Michalis-Menten (also called Holling type II) terms for substrate-limited processes such as nutrient uptake or grazing, Lotka-Volterra terms for some grazing functions, and linear mortality terms. The isoclines of these

- 1 functions are shown in Figure 1. We note that there exists a rich literature describing the
- 2 dependence of ecosystem model dynamics upon the forms of these terms. However, our
- 3 predator-prey subsystems contribute all the ecologically-realistic dynamics possible in
- 4 predator-prey systems, that is both stable spiral equilibrium points and stable limit cycles. We
- 5 could use other formulations to build our predator-prey subsystems, but this would not
- 6 contribute predator-prey subsystems with qualitatively different dynamics. Therefore, our
- 7 specific example reveals some quite general properties

9 The four equations of the *BFNPZ* model are:

10 
$$\frac{dB}{dt} = f_B B = \left[ \frac{k_1 (1 - k_{11})P}{P + k_2} + \frac{k_{25} (1 - k_{11})N}{N + k_{26}} - \frac{k_8 F}{B + k_9} - k_{10} \right] B, \tag{3}$$

11 
$$\frac{dF}{dt} = f_F F = \left[ \frac{k_8 (1 - k_{14})B}{B + k_9} - k_{13} \right] F, \tag{4}$$

12 
$$\frac{dP}{dt} = f_P P = \left[ \frac{k_{23}N}{N + k_{24}} - \frac{k_1 B}{P + k_2} - k_4 Z \right] P, \tag{5}$$

13 
$$\frac{dZ}{dt} = f_Z Z = \left[ k_4 \left( 1 - k_{20} \right) P - k_{19} \right] Z. \tag{6}$$

- 14 The closure of mass condition provides an extra model equation describing the concentration
- 15 of inorganic nutrient:

$$\frac{dN}{dt} = k_{10}B + k_{11} \left[ k_{25} \left( \frac{N}{N + k_{26}} \right) B + k_{1} \left( \frac{P}{P + k_{2}} \right) B \right] + k_{13}F + k_{8}k_{14} \left( \frac{B}{B + k_{9}} \right) F + k_{19}Z + k_{4}k_{20}PZ - k_{23} \left( \frac{N}{N + k_{24}} \right) P - k_{25} \left( \frac{N}{N + k_{26}} \right) B$$
(7)

- 1 As noted above, this equation (7) is not Kolmogorov form but is technically redundant as it
- 2 may be derived directly from the closure of mass condition:

$$N = N_T - B - F - P - Z \iff \frac{dN}{dt} = -\frac{dB}{dt} - \frac{dF}{dt} - \frac{dP}{dt} - \frac{dZ}{dt}, \tag{8}$$

- 4 where  $N_T$  is the total amount of nutrient contained in the system. Note that we could eliminate
- 5 N from equations (3) (6) by replacing it with  $N_T B F P Z$ , however, we will retain N
- 6 in the notation for simplicity. Then equations (3)-(6) are four equations in the four variables
- 7 BFPZ of Kolmogorov form, with 0 < B + F + P + Z < 1 when 0 < B, F, P, Z < 1 for all time.

- 9 Equations (3) (6) contain 16 parameters, values for which are based on field measurements
- that have been validated by simulating plankton dynamics in the Barents Sea (Gabric et al.
- 11 1999). The model is non-dimensionalised for our analysis; we used the maximum
- 12 phytoplankton growth rate  $k_{23}$  to define a characteristic time scale and define a total nutrient
- $N_T = 50$  to convert concentrations to proportions of the total. The measured parameter values
- are then replaced by their scaled equivalents:

15 
$$k'_m = \frac{k_m}{k_{23}}$$
 for  $m = 1, 8, 10, 13, 19, 23$  and 25, (9)

16 
$$k'_m = \frac{k_m}{N_T}$$
 for  $m = 2, 9, 24$  and 26, (10)

$$k_{4}' = \frac{k_{4}N_{T}}{k_{23}},\tag{11}$$

$$k'_{m} = k_{m}$$
 for  $m = 11,14$  and 20. (12)

1 The measured parameter values and their non-dimensional equivalents are given in Table 1.

2 The general constraints on valid parameter values for the model are that  $k_m > 0 \quad \forall m$  with the

slightly more restrictive constraint for the assimilation efficiency parameters that  $0 < k_m < 1$ 

for m = 11, 14 and 20. The model is therefore a Kolmogorov system, in that it may be written

5 in the form  $\dot{u}_i = f_i(u_1, u_2, K, u_n)u_i$ , i = 1, 2, K, n, and also meets Kolmogorov's *criteria* for

ecologically realistic dynamics of the BFN and NPZ predator-prey subsystems when

7 parameterised as above.

## 3. REDUCTION TO SUBSYSTEMS

The food web described by equations (3) - (6) is depicted in Figure 2 (top centre) with the autonomous food (sub-) webs that comprise it (Figure 2 centre and bottom rows). These subsystems may be obtained from the *BFNPZ* model by setting various state variables identically to zero in equations (3) - (6). The *BFNPZ* system, in that it is composed of the merged *BFNP* and *BNPZ* subsystems. The *BFNP* subsystem is in turn comprised of *BFN* and *BNP* subsystems, while the *BNPZ* subsystem is in turn comprised of *BPN* and *NPZ* subsystems. Diagrams of the trivial autotroph-nutrient systems *BN* and *NP* are not shown separately but are included in the two-variable subsystems. We similarly do not discuss the degenerate food webs (the *BNZ*, *FNP* and *FNZ* subsystems) that are technically possible but cannot survive in nature, and begin our considerations at the level of the three 'active' subsystems *BFN*, *BNP* and *NPZ*. We note that the autotrophs *B* and *P* are the 'drivers' of the ecosystem dynamics while the predators *F* and *Z* are 'passive' in that their oscillations and general dynamical behaviour follow that of their prey. Our claim will be that understanding the critical points, and their eigenvalues and eigenvectors, of these subsystems gives us significant

- 1 insight into the behaviour of the full system. In contrast, we require only of the fully interior
- 2 point of the full system that it be unstable (which it is for measured parameter values).

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

- 4 We present vector fields and long-term system trajectories for the BFNPZ model and
- 5 subsystems in Figure 3. This figure demonstrates two important properties of the model:
- that the dynamics of the models are mostly confined to near the boundaries
   (vertices, edges and faces) of the state spaces, and
  - and that the dynamics of each model can be inferred from the dynamics of its subsystems.

We reiterate that the BFN and NPZ models display the full spectrum of behaviours defined by Kolmogorov (1936) as possible for realistic predator-prey models: a stable limit cycle (BFN) and a stable spiral (NPZ). While we might explore different process representations in the models and generate different dynamics, the BFNPZ model contains all the dynamics that can realistically exist in ecosystem models under Kolmogorov's criteria. Although the vector fields in Figure 3 are specific to the BFNPZ model and its sub-models, and to the parameter values used in these models, the dynamics exhibited are general in that the orientation of the arrows of the vector fields in the regions of the spiral critical points is related to, and follows from, the Kolmogorov conditions that encapsulate much intuitive ecological thinking on appropriate predator-prey interaction behaviour. We would therefore expect similar behaviours to be exhibited by the models under parameter variations that maintained the validity of the Kolmogorov conditions, and the interaction omnivore behaviour, and kept the geometry of the fields similar to that shown. Note that the dynamics associated with the unstable predator-prey saddle point and the two stable autotroph nodes of the coupling (BNP) model, whilst a Kolmogorov system, do not comply with Kolmogorov's functional form criteria and hence do not exhibit predator-prey dynamics.

#### 4. CRITICAL POINTS

We now examine the critical points of the BFNPZ model and note that these are also critical points of the appropriate subsystems. We will therefore refer to each critical point in bold italics according to the system for which it is an 'internal' point, for example the critical point that has  $B, F, N \neq 0$  and P, Z = 0 will be denoted as the BFN critical point because it is internal in that system, but a boundary point in the BFNP and BFNPZ systems to which it also belongs. Note that because this nomenclature includes N for convenience, each critical point technically lies in a dimension one less than its name (i.e. the N point is a point of a zero dimension system at the origin of the Kolmogorov system, the BFN point is an interior point of a two-dimensional system, etc). Similarly, we report only the eigenvalues appropriate for the number of Kolmogorov dimensions for each point (i.e. the BFN point will have two eigenvalues in the BFN subsystem, but four in the full BFNPZ system).

We observe the heuristic that the  $f_i = 0$  of each 'feasible' subsystem (that is, not subsystems such as the BNZ, FNZ, FNP, BFNZ or FNPZ systems that cannot survive in nature) contribute a unique expression that identifies an interior point. This expression may have one or several roots, implying that the point may have simultaneously multiple locations in the state space. Often only one of these possibilities lies within the ecologically feasible region of the state space while the others lie outside. This is the case for all the critical points in our model system with the exception of the BNPZ point, which for the parameter set used, has only one root and this root lies outside the ecologically feasible state space. This point is described in the appendix but is not considered in the analysis as it does not influence the dynamics.

Analytic expressions for the critical points and their associated eigenvalues (where available)
are listed in the appendix with their numerical values calculated from the analytic expressions
for the parameter set in Table 1, or derived numerically if analytic expressions for the
eigenvalues are not available. The analytic expressions for the critical points and their
eigenvalues (where available) reveal the dependency on key parameters. A summary of the
critical point locations and their eigenvalues is given in Table 2.

- Table 2 reveals several striking features attesting to the consistency of the system with its subsystems:
  - (i) the critical points are consistent in each system, in that each system contains an internal critical point plus all the critical points of its subsystems
  - (ii) the eigenvalues (and hence the eigenvectors) of each critical point are the same in each system, that is knowledge of an eigenvalue of a point in a simple system allows that eigenvalue to be inferred for that point of the full system
  - (iii) the addition of extra species (dimensions) to a subsystem adds an eigenvalue to existing critical points that may be inferred from a simpler system, and adds a further internal critical point that may be similar to a subsystem critical point, but for which no information can be inferred a priori from the simpler systems
  - (iv) the eigenvalues of nearby critical points are similar, that is, the vector fields near the critical points are smooth.
  - We observe that features (iii) and (iv) together result in the complex eigenvalues of the *BFN* and *NPZ* predator-prey systems dominating as extra biota are added to form the higher dimension systems.

# 5. DYNAMICS

The consistencies of the critical points of the BFNPZ system and its subsystems leads to consistencies in the dynamical behaviours of the BFNPZ system and its subsystems (Figure 3). The BFN subsystem in Figure 3 (bottom left) exhibits a stable limit cycle, one of the two behaviours permitted under Kolmogorov's conditions. This subsystem includes the N saddle point at the origin, the BN autotroph point (a saddle in this subsystem) and the unstable spiral **BFN** predator-prey point (labelling as per Table 2). Note that the N and **BFN** critical points are located very near each other in the state space. In this case the unstable predator-prey critical point near the origin forces the orbits onto the stable limit cycle, the shape of which is determined by the eigenvectors of the other two critical points on the boundaries. The resultant

dynamics are the coexisting populations cycling forever.

The *NPZ* subsystem (Figure 3, bottom right) demonstrates a (spirally) stable critical point, the alternative behaviour permitted under Kolmogorov's conditions. Again this subsystem includes the *N* saddle point at the origin, the *NP* autotroph point (a saddle in this subsystem), and the stable spiral *NPZ* predator-prey point. This stable spiral point dominates the dynamics in the internal space of this subsystem, while the critical points on the boundaries direct the dynamics into the spiral. This behaviour is that of two species approaching an equilibrium (somewhat slowly in this case) by repeated, but decaying, oscillations.

The *BNP* subsystem (Figure 3 bottom centre) is not consistent with the Kolmogorov criteria (it remains however a Kolmogorov system) as the co-existence internal predator-prey point is not stable and the system must end up at one of the two stable autotroph points. This interesting

dynamic situation is due to *B* acting in this subsystem as an omnivore, that is, simultaneously both an autotroph feeding on *N* and a predator feeding on *P*. This system includes the *N* saddle point at the origin, the *BN* autotroph point, the *NP* autotroph point and a *BNP* predator-prey point. In this subsystem, the *NP* and *BNP* critical points are nearby. The interior *BNP* predator-prey point is a saddle and the two autotroph points (*BN* and *NP*) are both asymptotically stable. The dynamics of the system are therefore that either *B* or *P* dominates the subsystem. The separating surface that determines whether *B* or *P* dominates the subsystem

will be discussed below.

The dynamics of the *BFNP* subsystem (Figure 3 centre left) may be inferred by combining the dynamics of the *BFN* and *BNP* subsystems. This subsystem contains all the critical points (and eigenvalues) of the *BFN* and *BNP* subsystems (*N*, *BN*, *NP*, *BFN* and *BNP*) plus the further *BFNP* unstable spiral internal point. The critical points in this subsystem form groups, with the *N* and *BFN* points paired, and the *NP*, *BNP* and *BFNP* points forming a triple. The dynamics of the *BFNP* subsystem include the stable limit cycle dynamics of the *BFN* subsystem, but these successive *B*, *F* blooms are quickly overwhelmed by a *P* bloom that initiates when the system passes near the origin (the *N* saddle point) after which the subsystem is attracted along the axis to, and remains at, the stable *NP* autotroph point. Clearly, the separating surface near the origin that determines whether *B* or *P* blooms is critical to these dynamics.

Consideration of the dynamics of the *NPZ* and *BNP* subsystems similarly provides insight into the dynamics of the *BNPZ* subsystem (Figure 3 centre right). This subsystem contains all the critical points of the *NPZ* and *BNP* subsystems (*N*, *BN*, *NP*, *NPZ* and *BNP*) plus the further *BNPZ* saddle internal point (actually outside the state space in this example). Again the *N* and *BFN* points are paired, as are the *NP* and *BNP* points. In this case, the stable spiral of the *NPZ* 

subsystem is rendered unstable by the presence of B and instead of remaining at this point, the

system is initially attracted to it but is eventually directed away from it to the stable BN

3 autotroph point. In contrast to the BFNP subsystem, the dynamics in this subsystem occur

4 along the unstable eigenvector of the NPZ predator-prey point associated with B.

5

8

9

10

11

12

13

14

15

16

2

6 Finally, the dynamics of all the subsystems are evident in the full *BFNPZ* system (Figure 3

7 top). This system includes all the critical points of all the subsystems, although none of these

critical points are now stable. In particular, the autotroph critical points BN and NP that

formed the endpoints for the dynamics of the BFNP and BNPZ subsystems are now both

unstable because they both now have grazers feeding on them. Mathematically, the  $f_i$ 

associated with these grazers are positive at these points, and each contributes an unstable

eigenvector to the system. Therefore, rather than the dynamics starting with BFN dynamics and

ending at **NP** as in the BFNP subsystem, or starting in NPZ dynamics and finishing at **BN** as in

the BNPZ subsystem, the BFNPZ alternates between BFN and NPZ dynamics in a very robust

and stable limit cycle. The shape of this limit cycle is tightly controlled by the eigenvectors and

separating surfaces of the subsystems, as will be discussed below.

17

18

19

20

21

22

We also note that if we consider instead the reduction of the BFNPZ system to its subsystems,

we are confronted with somewhat counter-intuitive impacts of removing some species. For

example, if we remove F from the BFNPZ system to give the BNPZ system we effectively

initiate the extinction of the P and Z populations, as the BN critical point is now stable.

Similarly, if we remove Z from the BFNPZ system to give the BFNP system we effectively

initiate the extinction of the B and F populations, as the NP critical point is now stable.

24

#### **6. THE SIGNPOSTS**

2

1

3 The eigenvectors of the critical points on the boundaries of the state space of the BFNPZ 4 system, that is the critical points of the subsystems, provide 'signposts' that control the shape 5 of the limit cycle of the system. There are four such signposts; the N, BN, NP and NPZ critical 6 points. We observe that two of these critical points (N and NP) are located at vertices of the 7 state space, one (BN) is located on an edge and one (NPZ) is located on a face, clearly 8 explicating the role that the subsystems play in determining the full system dynamics. These 9 roles are revealed by the vector fields of the BFNPZ system in the vicinities of these points 10 (Figures 4 - 7); this will be further discussed below. 11 12 We also note that the one fully internal point of the system (the **BFNPZ** point) is important to 13 the dynamics in a different context, in that it (somewhat weakly) repels the system from the 14 interior of the state space so that the vector fields on the boundaries determine the dynamics. 15 Further, the **BFN** critical point associated with the BFN subsystem also plays a subtle role on a 16 face of the state space by similarly repelling trajectories and forcing the system to pass very 17 close to the *N* critical point.

18

19

20

21

22

23

24

The vector field at the origin of the state space (Figure 4) shows that trajectories will be attracted to the origin (the N critical point) and then repelled out along the unstable eigenvectors that lie along the B and P axes. (Note that the eigenvectors lie along the axes of the Kolmogorov state space but are shown in the non-Kolmogorov BNP state space.) The eigenvector along the P axis has an eigenvalue about 5 times that of the eigenvector along the B axis, indicating that P will grow faster than B in the region of the origin. Note also that the

1 **BFN** critical point has a repelling eigenvector, arising from the P competition, pointing into

the state space pushing trajectories towards the eigenvector along the P axis.

3

5

6

7

8

9

2

4 The vector field along the *P* axis from the origin maintains trajectories along the axis until this

eigenvector joins the corresponding attracting eigenvector, also lying along the P axis, at the

**NP** critical point (Figure 5). This is the largest eigenvalue of the 32 eigenvalues of the entire

system and is over double the magnitude of the next largest. This is therefore a very influential

point in determining the dynamics of the system. The stable eigenvectors of this point all lie

along axes of the Kolmogorov state space (the B, F and P axes). The unstable eigenvector

points inwards onto the PZ plane towards the NPZ critical point.

11

12

13

14

15

16

17

18

10

The imaginary eigenvectors of the spiral NPZ critical point (Figure 6) attract the trajectory

from the *NP* critical point while the repelling eigenvectors of the *BFNPZ* critical point and the

attracting real eigenvector of the **NPZ** critical point force the trajectory onto the **PZ** plane. The

trajectory only leaves this plane when it is pushed out along the unstable real eigenvector of the

**NPZ** critical point. This eigenvector arises from the B predation on P, and directs the system

through the interior of the state space towards the BN critical point. This is the only time the

limit cycle of *BFNPZ* system passes through the interior of the Kolmogorov state space.

19

20

21

22

23

24

The trajectory of the system is rapidly attracted across the state space by the large negative

eigenvalues of the BN critical point and then directed across the BF plane to the F axis by the

large positive eigenvector pointing in that direction. This eigenvector does not point to a

critical point, but the trajectories of the system are 'gathered' by the F axis and the vector field

indicates they are then directed along the F axis back to the origin where the cycle starts again.

- 1 The importance and subtlety of the origin in determining the dynamics of the full system, in
- 2 particular whether *B* or *P* blooms is revealed in the next section.

4

#### 7. SEPARATING SURFACE AT N CRITICAL POINT

6

5

7 The vector field of the BNP subsystem (Figure 8, main panel) reveals the general nature of this 8 'coupling' subsystem that links the BFN and NPZ subsystems. The sample trajectories shown 9 by the solid lines in Figure 8 reveal the influence of the strong negative eigenvalues of the 10 **BNP** critical point in drawing the trajectories to near the boundary of the state space where P dominates the system. The separatrix (dashed line), which separates trajectories that end up at 11 12 the BN critical point from those that end up at the NP critical point, suggests that most 13 trajectories of the system will end up at the BN critical point. In fact, for almost all cases (in 14 fact for 99% of the state space), B will eventually out-compete P and dominate the long-term 15 state of the system. This figure implies that most initial conditions will result in a B bloom 16 rather than a P bloom. However, the robust limit cycle we observe for the system, in which P 17 consistently blooms at the N critical point, and B consistently blooms only at the NPZ critical

point suggests that there is more to this story than is revealed by the main panel of Figure 8.

19

20

21

22

23

24

25

18

The region near the origin of Figure 8 is expanded in the inset, which also shows the separating surface (dashed line) that divides the region of the state space where the system will go to the BN critical point from the region that will go to the NP critical point. We observe that the fates of trajectories that pass near the origin of this state space are very different from those that do not, because the separatrix near the origin asymptotes to the B axis in the vicinity of B = 0.006. This reveals that for all concentrations of B lower than this level P will inevitably

dominate the system and bloom. The 'signature' dynamics of the system, with robustly

alternating P blooms and B breathers, is therefore critically dependant on the dynamics that

3 occur in the very small region of the state space close to the origin. The separating surface near

the N critical point therefore provides a useful insight into the scale of the interactions that are

important in understanding the dynamics of this system.

#### 8. BLOOMS AND BREATHERS

We now examine an interesting characteristic that arises from the coupling of the *BFN* system and the *NPZ* system. We will use the term "bloom" to describe the usual dynamics of *P* rapidly increasing from small levels to dominate the system, and introduce the term "breather" to describe the *B* outbreaks. This term has been associated with phenomena that emerge from exponentially small states to briefly dominate the system in partial differential equation models. The blooms of *B* that occur in the *BFNPZ* system for our measured parameter set can be described as breathers as they typically arise from extremely low concentrations. An interesting property of breathers is that they are effectively undetectable at almost all times; in a real ecosystem they would exist at levels that were unmeasurable or of the order of the measurement error. If the breather populations could be detected at times other than during an outbreak, their populations would be observed to be changing very slowly, and an outbreak could not be predicted from sparse knowledge of their changes in population over time.

To emphasise the bloom and breather dynamics, we now essentially decouple the dynamics of the BFN subsystem from those of the NPZ subsystem by calibrating the BFNPZ system with P parameter values that are typical of those used to fit P dynamics to Southern Ocean satellite

1 chlorophyll dynamics (Gabric et al. 2003). These parameter values result in an NPZ system

2 that is highly resilient (i.e. attracts to its NPZ spiral point very rapidly) and remains there for a

3 long time before B blooms. We then force this system with a sinusoidal cycle that emulates the

annual cycle of irradiance in the high latitudes of the Southern Ocean where phytoplankton

dynamics are closely related to the annual cycle of irradiance.

6

8

9

10

4

5

7 The resulting dynamics of the periodically forced system (Figure 9) reveal the presence of B

and F breathers every 9 - 13 years, interrupting the P and Z blooms that occur annually. The

system now exhibits somewhat benign chaos, where the timing and amplitude of the P, Z

blooms are mostly regular and predictable, but the B, F breathers are irregular and effectively

11 unpredictable.

12

13

14

15

16

17

18

19

20

21

22

23

An important implication for the capacity of systems such as the BFNPZ system to generate

breathers is a fundamental difference between numerical solvers for ordinary (ode) and partial

(pde) differential equations. While robust numerical solvers are available that can accurately

solve ode systems and resolve extremely small population sizes, the same cannot be said of

numerical solvers for pde systems. The constraints imposed by the requirements of resolving

spatial variation in the pde solvers make it difficult to resolve species concentrations to better

than one part in a million; whereas breathers typically require a resolution of at least one part in

a billion. We therefore observe that the endogenous dynamics that we observe in this analysis,

and which are an intrinsic property of the system equations, would not necessarily be observed

in models that resolve spatial variation. In particular, the breather dynamics that are evident

when populations recover from very low levels would not be reproduced in spatially resolved

models by pde solvers.

25

#### 9. DISCUSSION

This research has demonstrated the robust periodic dynamics of a moderate complexity plankton ecosystem model. These dynamics are comprised of regular 'blooms' of phytoplankton and zooplankton followed by 'breathers' of bacteria and zooflagellates that explode from exponentially small levels to briefly dominate the system. We examined the subsystems that comprise this system and noted counterintuitive behaviours. For example, removing the predator (F) of one prey (B) might reasonably be expected to permit the prey population to increase; however, it was not expected that such action would lead to the extinction of a whole related component of the food web (P and Z) leading to the effective collapse of the system. Similarly, removing Z from the system resulted in the dominance of P and the extinction of B and F.

The crucial point of the paper, however, is that we can understand the dynamics of the full system, the blooms and breathers, the robustly repeating cycles of the various individuals' struggle for life, as Rescigno and Richardson (1967) called it, in terms of the key two-variable subsystems. Our results indicate that the behaviour of a complex system can be inferred from the addition of the behaviours of its subsystems.

A key feature of our full dynamical system is that it is Kolmogorov in that each species change in time is proportional to its concentration in the sense that  $\dot{u}_i = f_i u_i$ . This property is automatically inherited by the subsystems when certain variables  $u_j \equiv 0$ . However, we note that we require further properties to achieve our realistic dynamics, that is, we need appropriate shapes or functional forms for the various  $f_i$ . May (1973) reiterated this point, first made by Kolmogorov (1936), and initiated a tradition of constraining ecosystem models to 'reasonable'

 $u_a \nabla f_a < 0 < u_p \nabla f_p$ , where a denotes an autotroph and p denotes a predator. Further conditions, which might be specified in terms of isocline intersections (Rescigno & Richardson 1967), are needed so that the key subsystems have only stable spiral attracting points or stable limit

functional forms by specifying various derivative conditions, of which we focus on

cycles, where the populations oscillate eternally. Our BFN and NPZ systems behave in this

manner, and we focus on the mass closure property to help render our full system in this

7 category.

It is expected that the instability of the interior critical point of the full system means that the dynamics can never rest in its neighbourhood. What is perhaps a little surprising is that other than this, our robust, stable periodic orbit appears not to be directly influenced by the eigenvalues and eigenvectors of this interior point. Instead, our orbit spends most of its time near vertices, edges and faces of the full system state space; i.e. near the N critical point (the origin, N = 1), the NP autotroph point (P = 1), the NPZ predator-prey point ( $P = P^*$ ,  $Z = Z^*$ ) and the BN autotroph point ( $B = B^*$ ). Physically, this dynamical systems behaviour manifests itself as successive blooms of P and Z interspersed with occasional breathers of B and F. Each population has its turn, even though each regularly decays to levels below  $10^{-6}$  of the total at some point in its life cycle. However, P and Z dominate the system most of the time.

The addition of an 'annual' forcing, analogous to the effects of seasonal changes in irradiance on phytoplankton, to the system adds a benign chaos to the system. Now, rather than appearing at regular intervals, the breathers become unpredictable, appearing every 9 - 13 'years' for our parameter values, chosen to be representative of Southern Ocean values. The precise magnitudes of the various breathers are similarly unpredictable, although the blooms occur with (reasonably) predictable magnitude at regular intervals. The regular bloom and breather

behaviour exhibited by the *BFNPZ* system exemplifies the resilience ideas of Holling (1973)

2 where the instability of interior critical points leads to highly resilient systems that are able to

3 robustly persist.

4

6

7

8

9

10

11

5 Finally, we note that the endogenous dynamics of the *BFNPZ* system that we have observed by

integrating the ordinary differential equations describing the biotic interactions will not

necessarily be evident when spatial variation is included and the partial differential equations

are solved. This is a result of intrinsic differences between ordinary and partial differential

equation solvers. We therefore suggest that when complex ecosystem models are coupled to,

for example, ocean circulation models, and when the timing and characteristics of bloom

events are critical, then particular care is taken to ensure that the numerical schemes correctly

12 handle small population levels.

13

#### 10. ACKNOWLEDGEMENTS

15

17

14

16 The authors wish to thank the Zilkha Trust, Lincoln College, Oxford for supporting this

research. We would also like to thank Fabio Boschetti and an anonymous reviewer for their

18 contributions to the manuscript.

19 20

#### 11. FIGURES AND TABLES

3

2

# 4 Figure Legends

5

- 6 Figure 1. Isoclines (upper panels) and function surfaces (lower panels) of the BFN (left), BNP
- 7 (middle) and NPZ (right) subsystems. The prey isoclines are shown as solid lines and the
- 8 predator isoclines as dotted lines in the upper panel.

9

- Figure 2. Food web of the BFNPZ model (top diagram) with the autonomous food webs that
- may be derived from it: BFNP model (middle left); BNPZ model (middle right); BFN model
- 12 (bottom left); BNP model (bottom centre) and NPZ model (bottom right).

13

- 14 Figure 3. Vector fields on the faces and dynamics of the BFNPZ (top), BFNP (middle left),
- 15 BNPZ (middle right), BFN (bottom left), BNP (bottom centre) and NPZ (bottom right) models.
- We have used N as a surrogate for F and Z in the BFNPZ model (top). When the dynamics are
- in the vicinity of the BN (left vertical) plane N effectively represents F, while when in the
- vicinity of the PN (right vertical) plane N effectively represents Z. The dynamics on the
- vertical planes in the top figure are therefore inverted from those of the lower figures.

20

21

- Figure 4. Critical points, vector field and eigenvectors in the vicinity of the N critical point of
- 23 the BFNPZ system. A part of the BFNPZ system limit cycle is also shown.

- Figure 5. Critical points, vector field and eigenvectors in the vicinity of the *P* critical point of
- 2 the BFNPZ system. A part of the BFNPZ system limit cycle is also shown.

- 4 Figure 6. Critical points, vector field and eigenvectors in the vicinity of the NPZ critical point
- of the *BFNPZ* system. A part of the *BFNPZ* system limit cycle is also shown.

6

- 7 Figure 7. Critical points, vector field and eigenvectors in the vicinity of the BN critical point of
- 8 the *BFNPZ* system. A part of the *BFNPZ* system limit cycle is also shown.

9

- Figure 8. Critical points, vector field and separating surface in the vicinity of the N, P and BNP
- critical points of the BFNPZ system. The dashed line marks the separating surface between
- initial conditions that go to the P critical point (below the dotted line) and those that go to the
- 13 BN critical point (above the dotted line). The solid lines are example trajectories of the system.
- 14 The region near the origin is blown up in the upper right, lines as for the larger figure.

15

- Figure 9. Blooms and breathers in the BFNPZ system. To create this figure,  $k_{13}$  was reduced to
- 17 0.05 to increase the time between blooms while  $k_{19}$  and  $k_{20}$  were varied to 0.50 and 0.38
- respectively to more rapidly dampen the P and Z oscillations. The P maximum growth term
- was forced with an equivalent annual cycle of irradiance  $k_{23}^{"} = k_{23} + 0.5 \sin(0.02\pi t)$ .

20

21

22

**Tables** 

23

24

Table 1: Parameter values used for the *BFNPZ* model.

PAR	PROCESS	UNITS	RAW	SCALED	REFERENCE
$k_I$	Max rate of N uptake by B	d <sup>-1</sup>	0.31	1.148	Muller-Niklas and Herndl (1996)
$k_2$	Half-sat const for B uptake of N	mgNm <sup>-3</sup>	34.65	0.693	Moloney et al. (1986)
$k_4$	Z grazing rate (per ind) on P	$m^3mgN^{-1}d^{-1}$	0.01	1.852	Gabric et al. (1999)
$k_8$	Max rate of B uptake by F	$d^{-1}$	1.67	6.185	Hansen et al. (1996)
$k_9$	Half-sat const for F uptake of B	mgNm <sup>-3</sup>	9.10	0.182	Fenchel (1982)
$k_{I0}$	B specific excretion rate	$d^{-1}$	0.07	0.259	Moloney et al. (1986)
$k_{II}$	Prop of N uptake excreted by B	-	0.63	0.63	Moloney et al. (1986)
$k_{13}$	F specific excretion rate	$d^{-1}$	0.05	0.1851	Moloney et al. (1986)
$k_{14}$	Prop of N uptake excreted by F	-	0.65	0.65	Moloney et al. (1986)
$k_{19}$	Z specific N excretion rate	$d^{-1}$	0.05	0.1852	Moloney et al. (1986)
$k_{20}$	Prop of N uptake excreted by Z	-	0.40	0.40	Moloney et al. (1986)
$k_{23}$	Max rate of N uptake by P	$d^{-1}$	0.27	1	Gabric et al. (1999)
$k_{24}$	Half-sat const for P uptake of N	mgNm <sup>-3</sup>	12.60	0.252	Slagstad and Stole-Hansen (1991)
$k_{25}$	Max rate of N uptake by B	$d^{-1}$	0.31	1.148	Muller-Niklas and Herndl (1996)
$k_{26}$	Half-sat const for B uptake of N	mgNm <sup>-3</sup>	3.45	0.069	Billen and Becquevort (1991)
$N_T$	Total nutrient as nitrogen	mgNm <sup>-3</sup>	50	1	Gabric et al. (1999)

Table 2. Critical point values and associated eigenvalues.

СР	SYS	<b>B</b> *	$F^*$	$N^*$	<b>P</b> *	$Z^*$	$\lambda_{_{1}}$	$\lambda_2$	$\lambda_3$	$\lambda_4$	ST <sup>1</sup>
	BN	0	-	1	-	-	0.1383	-	-	-	US
	N <b>P</b>	-	-	1	0	-	-	-	0.7987	-	US
	<b>BFN</b>	0	0	1			0.1383	-0.1851	-	-	SAD
<b>3</b> .7	BNP	0	-	1	0	-	0.1383	-	0.7987	-	US
N	<b>NPZ</b>	-	-	1	0	0	-	-	0.7987	-0.1852	SAD
	BFNP	0	0	1	0	-	0.1383	-0.1851	0.7987	-	SAD
	BNPZ	0	-	1	0	0	0.1383	-	0.7987	-0.1852	SAD
	BFNPZ	0	0	1	0	0	0.1383	-0.1851	0.7987	-0.1852	SAD
	BN	0.8919	-	0.1081		_	-0.8364	-	-	-	AS
	<b>BFN</b>	0.8919	0	0.1081	-	-	-0.8364	1.6129	-	-	SAD
<b>D</b> 1/	BNP	0.8919	-	0.1081	0	-	-0.8364	-	-1.1783	-	AS
BN	BFNP	0.8919	0	0.1081	0	-	-0.8364	1.6129	-1.1783	-	SAD
	BNPZ	0.8919	-	0.1081	0	0	-0.8364	-	-1.1783	-0.1852	AS
	BFNPZ	0.8919	0	0.1081	0	0	-0.8364	1.6129	-1.1783	-0.1852	SAD
	N <b>P</b>	-	-	0	1	-	-	-	-3.9683	-	AS
	BNP	0	-	0	1	-	-0.0081	-	-3.9683	-	AS
	NPZ	-	-	0	1	0	-	-	-3.9683	0.9260	SAD
NP	BFNP	0	0	0	1	0	-0.0081	-0.1851	-3.9683	-	AS
	BNPZ	0	0	0	1	0	-0.0081	-	-3.9683	0.9260	SAD
	BFNPZ	0	0	0	1	0	-0.0081	-0.1851	-3.9683	0.9260	SAD
	BFN	0.0170	0.0044	0.9785	-	-	0.0057 ±	0.1527 <i>i</i>	-	-	US
<b>BFN</b>	BFNP	0.0170	0.0044	0.9785	0	-	0.0057 ±	0.1527 <i>i</i>	0.7670	-	US
	BFNPZ	0.0170	0.0044	0.9785	0	0	0.0057 ±	0.1527 <i>i</i>	0.7670	-0.1852	SSAD
<b>D</b> 37**	BNP	0.0091	-	0.0016	0.9894	-	0.0079	-	-3.9365	-	SAD
BNP	BNPZ	0.0091	-	0.0016	0.9894	0	0.0079	_	-3.9365	0.9145	SAD

	BFNPZ	0.0091	0	0.0016	0.9894	0	0.0079	-0.0852	-3.9365	0.9145	SAD
	NPZ	-	-	0.4794	0.1667	0.3539	-	-	-0.0393	± 0.3882i	SS
NPZ	BNPZ	0	-	0.4794	0.1667	0.3539	0.1947	-	$-0.0393 \pm 0.3882i$		US
	BFNPZ	0	0	0.4794	0.1667	0.3539	0.1947	-0.1851	-0.0393	± 0.3882 <i>i</i>	SSAD
$BFNP^2$	BFNP	0.0170	0.0002	0.0030	0.9798	-	$0.0076 \pm 0.0340i$		-3.9017	-	SSAD
DFNF	BFNPZ	0.0170	0.0002	0.0030	0.9798	0	$0.0076 \pm 0.0340i$		-3.9017	0.9037	SSAD
BFNPZ <sup>2</sup>	BFNPZ	0.0170	0.0062	0.4707	0.1667	0.3394	$0.0079 \pm 0.1799i$		$-0.0384 \pm 0.3839i$		SSAD

<sup>&</sup>lt;sup>1</sup> SAD denotes a saddle point (unstable in at least one dimension), SSAD denotes a spiral saddle point, AS denotes an asymptotically stable point, SS denotes a spirally stable point and US denotes an unstable node.

<sup>2</sup> Eigenvalues were calculated numerically for the *BFNP* and *BFNPZ* points except for  $\lambda_4$  of the *BFNP* point for which an analytic expression was obtained (see appendix).

#### 12. LITERATURE CITED

- Anderson TR (2005) Plankton functional type modelling: running before we can walk? J Plankton Res 27:1073-1081
- Arhonditsis GB, Brett MT (2004) Evaluation of the current state of mechanistic aquatic biogeochemical modeling. Marine Ecology Progress Series 271:13-26
- Baretta J, Ebenhoh W, Ruardij P (1995) The European Regional Seas Ecosystem Model, a complex marine ecosystem model. Journal of Sea Research 33:233-246
- Charlson RJ, Lovelock JE, Andreae MO, Warren SG (1987) Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. Nature 326:655-661
- Cropp RA, Norbury J, Gabric AJ, R. D. Braddock (2004), Global Biogeochem. Cycles, 18, (2004) Modeling dimethylsulphide production in the upper ocean. Global Biogeochemical Cycles 18:doi:10.1029/2003GB002126.
- Dugdale RC, Goering JJ (1967) Uptake of new and regenerated forms of nitrogen in primary productivity. Limnology and Oceanography 12:196-206
- Edwards AM, Brindley J (1999) Zooplankton mortality and the dynamical behaviour of plankton population models. Bulletin Of Mathematical Biology 61:303-339
- Falkowski P, Scholes RJ, Boyle E, Canadell J, Canfield D, Elser J, Gruber N, Hibbard K, Hogberg P, Linder S, Mackenzie FT, Moore B, Pedersen T, Rosenthal Y, Seitzinger S, Smetacek V, Steffen W (2000) The global carbon cycle: A test of our knowledge of the Earth as a system. Science 290:291-296
- Franks PJS (2002) NPZ models of plankton dynamics: their construction, coupling to physics, and application. Journal of Oceanography 58:379-387
- Gabric AJ, Cropp RA, Hirst AC, Marchant HJ (2003) The response of dimethylsulphide production to simulated warming in the eastern Antarctic Southern Ocean. Tellus 55B:966-981
- Gabric AJ, Matrai PA, Vernet M (1999) Modelling the production and cycling of dimethylsulphide during the vernal bloom in the Barents Sea. Tellus 51B:919-937
- Gibson G, Musgrave D, Hinckley S (2005) Non-linear dynamics of a pelagic ecosystem model with multiple predator and prey types. Journal of Plankton Research 27:427-447
- Holling CS (1973) Resilience and stability of ecological systems. Annual Review of Ecology and Systematics 4:1-23
- Huang XC, Zhu L (2005) Limit cycles in a general Kolmogorov model. Nonlinear Analysis 60:1393-1414
- Kolmogorov AN (1936) Sulla Teoria di Volterra della Lotta per l'Esisttenza. Giorn Instituto Ital Attuari 7:74-80
- Kuznetsov Y, Rinaldi S (1996) Remarks on Food Chain Dynamics. Mathematical Biosciences 134:1-33
- Lancelot C, Hannon E, Becquevort S, Veth C, De Baar HJW (2000) Modeling phytoplankton blooms and carbon export production in the Southern Ocean: dominant controls by light and iron in the Atlantic sector in Austral spring 1992. Deep-Sea Research I 47:1621-1662
- Le Quere C, Harrison SP, Prentice IC, Buitenhuis ET, Aumonts O, Bopp L, Claustre H, Cotrim da Cunha L, Geider RJ, Giraud X, Klaas C, Kohfeld KE, Legrende L, Manizza M, Platt T, Rivkin RB, Sathyendranath S, Uitz J, Watson A, Wolf-Gladrow D (2005) Ecosystem

- dynamics based on plankton functional types for global ocean biogeochemistry models. Global Change Biology 11:2016-2040
- Matear RJ (1995) Parameter optimization and analysis of ecosystem models using simulated annealing: A case study at Station P. Journal of Marine Research 53:571-607
- May RM (1973) Stability and Complexity in Model Ecosystems., Vol. Princeton University Press, Princeton
- Moloney CL, Bergh MO, Field JG, Newell RC (1986) The effect of sedimentation and microbial nitrogen regeneration in a plankton community: a simulation investigation. Journal of Plankton Research 8:427-445
- Neutel A-M, Heesterbeek JAP, de Ruiter PC (2002) Stability in real food webs: Weak links in long loops. Science 296:1120-1123
- Neutel A-M, Heesterbeek JAP, van de Koppel J, Hoenderboom G, Vos A, Kaldeway C, Berendse F, de Ruiter PC (2007) Reconciling complexity with stability in naturally assembling food webs. Nature 449:599-602
- Rescigno A, Richardson IW (1967) The Struggle for Life I. Two Species. Bull Mathematical Biophysics 29:377-388
- Seppelt R, Richter O (2005) "It was an artefact not the result": A note on systems dynamic model development tools. Environmental Modelling And Software 20:1543-1548
- Spitz YH, Moisan JR, Abbott MR (2001) Configuring an ecosystem model using data from the Bermuda Atlantic Time Series (BATS). Deep-Sea Research II 48:1733-1768
- Vallina SM, Simo R, Popova EE, Anderson TR, Gabric A, Cropp RA, Pacheco JM (2008) A dynamic model of ocean sulfur (DMOS) applied to the Sargasso Sea: simulating the dimethylsulfide (DMS) summer paradox. Journal of Geophysical Research (Biogeosciences) 113:G01009, doi:01010.01029/02007JG000415

#### 13. APPENDIX

# N critical point

In common with all Kolmogorov models, the *BFNPZ* model has a critical point at the origin of the system where no biota are extant:

$$N_N^* = 1, (13)$$

$$B_N^*, F_N^*, P_N^*, Z_N^* = 0. (14)$$

The eigenvalues of the Jacobian at this point are:

$$\lambda_{N-1} = \frac{k_{25} \left( 1 - k_{11} \right)}{1 + k_{26}} - k_{10} = 0.1383, \tag{15}$$

$$\lambda_{N-2} = -k_{13} = -0.1851, \tag{16}$$

$$\lambda_{N-3} = \frac{k_{23}}{1 + k_{24}} = 0.7987, \tag{17}$$

$$\lambda_{N-4} = -k_{19} = -0.1852. \tag{18}$$

The eigenvectors associated with these eigenvalues all point along the species axes, with the unstable autotroph eigenvectors associated with  $\lambda_{N-1}$  and  $\lambda_{N-3}$  pointing along the B and P axes respectively.

#### **BN** critical point

The first autotroph critical point of the *BFNPZ* model is given by:

$$B_{BN}^* = 1 - \frac{k_{10}k_{26}}{k_{25}(1 - k_{11}) - k_{10}} = 0.8919,$$
 (19)

$$N_{BN}^* = \frac{k_{10}k_{26}}{k_{25}(1 - k_{11}) - k_{10}} = 0.1081, \tag{20}$$

$$F_{BN}^*, P_{BN}^*, Z_{BN}^* = 0. (21)$$

The eigenvalues of this point are:

$$\lambda_{BN-1} = k_{25} \left( 1 - k_{11} \right) \left( \frac{N_{BN}^{*-2} + k_{26} \left( N_{BN}^{*} - B_{BN}^{*} \right)}{\left( N_{BN}^{*} + k_{26} \right)^{2}} \right) - k_{10} = -0.8364, \quad (22)$$

$$\lambda_{BN-2} = k_8 \left( 1 - k_{14} \right) \left( \frac{B_{BN}^*}{B_{BN}^* + k_9} \right) - k_{13} = 1.6129, \tag{23}$$

$$\lambda_{BN-3} = k_{23} \left( \frac{N_{BN}^*}{N_{BN}^* + k_{24}} \right) - \left( \frac{k_1}{k_2} \right) B_{BN}^* = -1.1783, \tag{24}$$

$$\lambda_{RN-4} = -k_{19} = -0.1852. \tag{25}$$

The eigenvectors associated with these eigenvalues again all point along the species axes, with the exception of the unstable eigenvector associated with  $\lambda_{BN-2}$  which points internally towards the F axis. Note that the  $\lambda_{N-1}$  eigenvector pointing away from the N critical point

joins the  $\lambda_{BN-1}$  eigenvector pointing into the **BN** critical point. A B bloom initiated at the origin must therefore end up at the **BN** critical point.

# **NP** critical point

The second autotroph-only critical point of the model is:

$$P_{NP}^* = 1$$
, (26)

$$B_{NP}^*, F_{NP}^*, N_{NP}^*, Z_{NP}^* = 0. (27)$$

The eigenvalues of this point are:

$$\lambda_{NP-1} = \frac{k_1 \left(1 - k_{11}\right)}{1 + k_2} - k_{10} = -0.0081, \tag{28}$$

$$\lambda_{NP-2} = -k_{13} = -0.1851, \tag{29}$$

$$\lambda_{NP-3} = -\frac{k_{23}}{k_{24}} = -3.9683, \tag{30}$$

$$\lambda_{NP-4} = k_4 \left( 1 - k_{20} \right) - k_{19} = 0.9620$$
 (31)

Once again, the eigenvectors associated with these eigenvalues again all point along the species axes, with the exception of the unstable eigenvector associated with  $\lambda_{NP-4}$  which points along a separatrix into the NPZ plane. Note in this case that the  $\lambda_{N-3}$  eigenvector pointing away from the N critical point joins the  $\lambda_{NP-3}$  eigenvector pointing into the NP critical point. A P bloom initiated at the origin must therefore end up at the NP critical point.

#### **BFN** critical point

The first predator-prey critical point of the BFNPZ model is at:

$$B_{BFN}^* = \frac{k_9 k_{13}}{k_8 \left(1 - k_{14}\right) - k_{13}} = 0.0170,$$
(32)

$$F_{BFN}^* = \frac{k_{25}}{k_8} \left( 1 - k_{11} \right) \left( B_{BFN}^* + k_9 \right) \left( \frac{N_{BFN}^*}{N_{BFN}^* + k_{26}} \right) - \frac{k_{10}}{k_8} \left( B_{BFN}^* + k_9 \right) = 0.0044, (33)$$

$$N_{BFN}^{*} = \frac{1}{2} \begin{bmatrix} -\left(k_{26} + B_{BFN}^{*} + \left(\frac{B_{BFN}^{*} + k_{9}}{k_{8}}\right) \left(k_{25} \left(1 - k_{11}\right) - k_{10}\right) - 1\right) \\ \left(k_{26} + B_{BFN}^{*} + \left(\frac{B_{BFN}^{*} + k_{9}}{k_{8}}\right) \left(k_{25} \left(1 - k_{11}\right) - k_{10}\right) - 1\right)^{2} \\ -4k_{26} \left(B_{BFN}^{*} - 1 + k_{10} \left(\frac{B_{BFN}^{*} + k_{9}}{k_{8}}\right)\right) \end{bmatrix} = 0.9785, \quad (34)$$

$$P_{BFN}^*, Z_{BFN}^* = 0. (35)$$

The eigenvalues at this point are:

$$\operatorname{Re}\left(\lambda_{BFN-1,2}\right) = k_{8} \left(\frac{B_{BFN}^{*}}{\left(B_{BFN}^{*} + k_{9}\right)^{2}}\right) - k_{25} \left(1 - k_{11}\right) \left(\frac{k_{26}}{\left(N_{BFN}^{*} + k_{26}\right)^{2}}\right) B_{BFN}^{*} = 0.0057, \quad (36)$$

$$\begin{bmatrix}
k_{8} \left( \frac{B_{BFN}^{*}}{\left( B_{BFN}^{*} + k_{9} \right)^{2}} \right) F_{BFN}^{*} - k_{25} \left( 1 - k_{11} \right) \left( \frac{k_{26}}{\left( N_{BFN}^{*} + k_{26} \right)^{2}} \right) B_{BFN}^{*} \right]^{2} - \\
4 \left[ k_{25} \left( 1 - k_{11} \right) \left( \frac{k_{26}}{\left( N_{BFN}^{*} + k_{26} \right)^{2}} \right) B_{BFN}^{*} + k_{8} \left( \frac{B_{BFN}^{*}}{B_{BFN}^{*} + k_{9}} \right) \right] \times \\
\left[ k_{8} \left( 1 - k_{14} \right) \left( \frac{k_{9}}{\left( B_{BFN}^{*} + k_{9} \right)^{2}} \right) F_{BFN}^{*} \right] \\
= \pm 0.1527i.$$
(37)

$$\lambda_{BFN-3} = k_{23} \left( \frac{N_{BFN}^*}{N_{BFN}^* + k_{24}} \right) - \left( \frac{k_1}{k_2} \right) B_{BFN}^* = 0.7670,$$
 (38)

$$\lambda_{BFN-4} = -k_{19} = -0.1852 \,. \tag{39}$$

The unstable eigenvectors of this point push orbits towards the origin, from where the orbit is directed to the *NP* autotroph critical point.

### **BNP** critical point

The second predator-prey critical point (technically an omnivore-prey critical point as B consumes both N and P) is given by:

$$B_{BNP}^* = \frac{k_{23} \left( P_{BNP}^* + k_2 \right) N_{BNP}^*}{k_1 \left( N_{BNP}^* + k_{24} \right)} = 0.0091, \tag{40}$$

$$\alpha N_{BNP}^{*3} + \beta N_{BNP}^{*2} + \chi N_{BNP}^{*} + \delta = 0 \quad \Rightarrow N_{BNP}^{*} = 0.0016, \tag{41}$$

$$P_{BNP}^{*} = \frac{k_{2} \left[ k_{10} \left( N_{BNP}^{*} + k_{26} \right) - k_{25} \left( 1 - k_{11} \right) N_{BNP}^{*} \right]}{k_{1} \left( 1 - k_{11} \right) \left( N_{BNP}^{*} + k_{26} \right) + k_{25} \left( 1 - k_{11} \right) N_{BNP}^{*} - k_{10} \left( N_{BNP}^{*} + k_{26} \right)} = 0.9894, \tag{42}$$

$$F_{BNP}^*, Z_{BNP}^* = 0, (43)$$

where:  $\alpha = k_1 \left[ k_1 + k_{25} - k_{10} - k_{11} \left( k_1 + k_{25} \right) \right]$ 

$$\beta = k_1 \begin{bmatrix} k_{11} (k_{25} \{1 - k_{24} + k_2\} + k_1 \{1 - k_{24} - k_{26}\} - k_2 k_{23}) + k_1 (k_{24} + k_{26} - 1) \\ -k_{10} (k_{26} - k_2 + k_{24} - 1) + k_2 k_{23} - k_{25} (k_2 - k_{24} + 1) \end{bmatrix},$$

$$\chi = k_{1} \begin{bmatrix} k_{1} \left( k_{26} \left\{ k_{24} - 1 + k_{11} \right\} - k_{24} + k_{11} k_{24} \left\{ 1 - k_{26} \right\} \right) \\ + k_{24} k_{25} \left( 1 + k_{2} \right) \left( k_{11} - 1 \right) + k_{10} k_{26} \left( 1 + k_{2} \right) \\ + k_{2} k_{23} k_{26} \left( 1 - k_{11} \right) + k_{10} k_{24} \left( 1 + k_{2} - k_{26} \right) \end{bmatrix},$$

$$\delta = k_1 \left[ k_{24} k_{26} \left( k_{10} \left\{ k_2 + 1 \right\} - k_1 \left\{ 1 - k_{11} \right\} \right) \right].$$

The eigenvalues at this point are:

$$\lambda_{BNP-1} = \frac{\alpha' + \beta' + \sqrt{(\alpha' + \beta')^2 + 4\chi'\delta'}}{2} = 0.0079,$$
 (44)

$$\lambda_{BNP-2} = k_8 \left( 1 - k_{14} \right) \left( \frac{B_{BNP}^*}{B_{BNP}^* + k_9} \right) - k_{13} = -0.0852, \tag{45}$$

$$\lambda_{BNP-3} = \frac{\alpha' + \beta' - \sqrt{(\alpha' + \beta')^2 + 4\chi'\delta'}}{2} = -3.9365,$$
 (46)

$$\lambda_{BNP-4} = k_4 \left( 1 - k_{20} \right) P_{BNP}^* - k_{19} = 0.9145 \,, \tag{47}$$

where

$$\alpha' = k_{1} \left( 1 - k_{11} \right) \left( \frac{P_{BNP}^{*}}{P_{BNP}^{*} + k_{2}} \right) + k_{25} \left( 1 - k_{11} \right) \left( \frac{N_{BNP}^{*}}{N_{BNP}^{*} + k_{26}} \right)$$

$$-k_{25} \left( 1 - k_{11} \right) \left( \frac{k_{26}}{\left( N_{BNP}^{*} + k_{26} \right)^{2}} \right) B_{BNP}^{*} - k_{10},$$

$$(48)$$

$$\beta' = k_{23} \left( \frac{N_{BNP}^*}{N_{BNP}^* + k_{24}} \right) - k_{23} \left( \frac{k_{24}}{\left(N_{BNP}^* + k_{24}\right)^2} \right) P_{BNP}^* - k_1 \left( \frac{k_2}{\left(P_{BNP}^* + k_2\right)^2} \right) B_{BNP}^*, (49)$$

$$\chi' = -k_{23} \left( \frac{k_{24}}{\left( N_{BNP}^* + k_{24}^* \right)^2} \right) P_{BNP}^* - k_1 \left( \frac{P_{BNP}^*}{P_{BNP}^* + k_2} \right), \tag{50}$$

$$\delta' = -k_1 \left( 1 - k_{11} \right) \left( \frac{k_2}{\left( P_{BNP}^* + k_2 \right)^2} \right) B_{BNP}^*. \tag{51}$$

#### **NPZ** critical point

The third predator-prey critical point of the model is given by:

$$B_{NPZ}^*, F_{NPZ}^* = 0, (52)$$

$$N_{NPZ}^{*} = \frac{1}{2} \left[ -\left( k_{24} + P_{NPZ}^{*} + \frac{k_{23}}{k_{4}} - 1 \right) \pm \sqrt{\left( k_{24} + P_{NPZ}^{*} + \frac{k_{23}}{k_{4}} - 1 \right)^{2} - 4k_{24} \left( P_{NPZ}^{*} - 1 \right)} \right], \quad (53)$$

$$= 0.4794$$

$$P_{NPZ}^* = \frac{k_{19}}{k_4 \left( 1 - k_{20} \right)} = 0.1667 \,, \tag{54}$$

$$Z_{NPZ}^* = \frac{k_{23}}{k_4} \left( \frac{N_{NPZ}^*}{N_{NPZ}^* + k_{24}} \right) = 0.3539.$$
 (55)

The eigenvalues at this point are:

$$\lambda_{NPZ-1} = k_1 \left( 1 - k_{11} \right) \left( \frac{P_{NPZ}^*}{P_{NPZ}^* + k_2} \right) + k_{25} \left( 1 - k_{11} \right) \left( \frac{N_{NPZ}^*}{N_{NPZ}^* + k_{26}} \right) - k_{10} = 0.1947, (56)$$

$$\lambda_{NPZ-2} = -k_{13} = -0.1851, \tag{57}$$

$$\operatorname{Re}\left(\lambda_{NPZ-3,4}\right) = -\frac{k_{23}P_{NPZ}^*}{2} \left[ \frac{k_{24}}{\left(N_{NPZ}^* + k_{24}\right)^2} \right] = -0.0393, \tag{58}$$

$$\operatorname{Im}(\lambda_{NPZ-3,4}) = \pm \frac{1}{2} \sqrt{\frac{\left\{k_{23}P_{NPZ}^* \left[\frac{k_{24}}{\left(N_{NPZ}^* + k_{24}\right)^2}\right]\right\}^2}{+4k_4\left(1 - k_{20}\right)\left\{k_{23}\left[\frac{k_{24}}{\left(N_{NPZ}^* + k_{24}\right)^2}\right] - k_4\right\}P_{NPZ}^*Z_{NPZ}^*}} = \pm 0.3882i. (59)$$

The unstable eigenvector associated with  $\lambda_{NPZ-1}$  points toward the **BN** autotroph critical point.

#### **BFNP** critical point

The first higher-dimension food web critical point, where four of the five state variables are non-zero, is given by:

$$B_{BFNP}^* = \frac{k_9 k_{13}}{k_8 \left(1 - k_{14}\right) - k_{13}} = 0.0170, \tag{60}$$

$$F_{BFNP}^* = \frac{k_{25}}{k_8} \left( 1 - k_{11} \right) \left( B_{BFNP}^* + k_9 \right) \left( \frac{N_{BFNP}^*}{N_{BFNP}^* + k_{26}} \right) - \frac{k_{10}}{k_8} \left( B_{BFNP}^* + k_9 \right) = 0.0002, \tag{61}$$

$$N_{BFNP}^{*}{}^{4} + N_{BFNP}^{*}{}^{3} \left[ k_{26} + k_{24} - 1 - k_{2} + B_{BFNP}^{*} - \beta k_{10} + \alpha + \beta \gamma \left( 1 - \frac{k_{2}}{\alpha} \right) + \beta \delta \right]$$

$$+ N_{BFNP}^{*}{}^{2} \left[ k_{26} \left( k_{24} - 1 - k_{2} + B_{BFNP}^{*} - \beta k_{10} + \alpha + \beta \gamma \left( 1 - \frac{k_{2}}{\alpha} \right) \right) + \alpha k_{24} \right]$$

$$- k_{24} \left( 1 + k_{2} - B_{BFNP}^{*} + \beta k_{10} - \alpha \right) + \beta k_{24} \left( \delta + \gamma \right)$$

$$+ N_{BFNP}^{*} \left[ \alpha k_{24}^{2} + k_{26} \left( \alpha k_{24} - k_{24} \left[ 1 + k_{2} - B_{BFNP}^{*} + \beta k_{10} - \alpha \right] + \beta \gamma k_{24} \right) \right] + \alpha k_{24}^{2} k_{26} = 0$$

$$\Rightarrow N_{BFNP}^{*} = 0.0030,$$

$$(62)$$

$$P_{BFNP}^* = \frac{k_1}{k_{23}} \left( \frac{N + k_{24}}{N} \right) B_{BFNP}^* - k_2 = 0.9798, \tag{63}$$

$$Z_{BFNP}^* = 0, (64)$$

where 
$$\alpha = \frac{k_1 B_{BFNP}^*}{k_{23}}$$
,  $\beta = \frac{B_{BFNP}^* + k_9}{k_8}$ ,  $\delta = k_{25} \left(1 - k_{11}\right)$ ,  $\varepsilon = \frac{k_2 k_{23}}{k_1 B_{BFNP}^*}$  and  $\gamma = k_1 \left(1 - k_{11}\right)$  and only

the positive roots are ecologically feasible.

Analytic expressions for most of the eigenvalues at the BFNP point are not useful, and these eigenvalues will be calculated numerically. However, the analytic expression for the eigenvalue associated with the Z dimension is informative:

$$\lambda_{BFNP-4} = k_4 \left( 1 - k_{20} \right) P_{BFNP}^* - k_{19} = 0.9037. \tag{65}$$

We note from comparison with the numerically calculated eigenvalues listed in Table 3 that the  $\lambda_{BFNP-4}$  eigenvalue is the most unstable direction at this critical point, by a factor of about 100.

### BNPZ critical point

The second higher-dimension food web critical point, where four of the five state variables are non-zero, is given by:

$$B_{BNPZ}^{*} = \frac{\left(P_{BNPZ}^{*} + k_{2}\right)\left[1 - N_{BNPZ}^{*} - P_{BNPZ}^{*} - \frac{k_{23}}{k_{4}}\left(\frac{N_{BNPZ}^{*}}{N_{BNPZ}^{*} + k_{24}}\right)\right]}{P_{BNPZ}^{*} + k_{2} - \frac{k_{1}}{k_{4}}} = 2.4959, (66)$$

$$F_{BNPZ}^* = 0, (67)$$

$$N_{BNPZ}^{*} = \frac{k_{26} \left[ k_{1} \left( 1 - k_{11} \right) P_{BNPZ}^{*} - k_{10} \left( P_{BNPZ}^{*} + k_{2} \right) \right]}{\left( P_{BNPZ}^{*} + k_{2} \right) \left[ k_{10} - k_{25} \left( 1 - k_{11} \right) \right] - k_{1} \left( 1 - k_{11} \right) P_{BNPZ}^{*}} = 0.0492, \quad (68)$$

$$P_{BNPZ}^* = \frac{k_{19}}{k_4 \left( 1 - k_{20} \right)} = 0.1667 \,, \tag{69}$$

$$Z_{BNPZ}^* = \frac{k_{23}}{k_4} \left( \frac{N_{BNPZ}^*}{N_{BNPZ}^* + k_{24}} \right) - \frac{k_1}{k_4} \left( \frac{B_{BNPZ}^*}{P_{BNPZ}^* + k_2} \right) = -1.7118.$$
 (70)

Analytic expressions for most of the eigenvalues at the BNPZ point are not useful, and these eigenvalues will be calculated numerically. However, the analytic expression for the eigenvalue associated with the F dimension is informative:

$$\lambda_{BNPZ-2} = k_8 \left( 1 - k_{14} \right) \left( \frac{B_{BNPZ}^*}{B_{BNPZ}^* + k_9} \right) - k_{13} = 1.8326.$$
 (71)

Note that for the parameter values used in the model, this point lies outside the ecologically feasible region of the state space, and the eigenvalues do not have any influence on the dynamics of the system. We further note that this point may be brought into the feasible state space with subtle variations of certain parameter values.

## BFNPZ critical point

The final critical point of the *BFNPZ* model, that is not shared by any of the sub-models, the only critical point to have all positive, non-zero state variables (i.e. the only 'interior' point of the full model) is:

$$B_{BFNPZ}^* = \frac{k_9 k_{13}}{k_8 \left(1 - k_{14}\right) - k_{13}} = 0.0170,$$
 (72)

$$F_{BFNPZ}^* = \left(\frac{B_{BFNPZ}^* + k_9}{k_8}\right) \left[k_1 \left(1 - k_{11}\right) \left(\frac{P_{BFNPZ}^*}{P_{BFNPZ}^* + k_2}\right) + k_{25} \left(1 - k_{11}\right) \left(\frac{N_{BFNPZ}^*}{N_{BFNPZ}^* + k_{26}}\right) - k_{10}\right] (73)$$

$$= 0.0062,$$

$$N_{BFNPZ}^{*}^{3} + N_{BFNPZ}^{*}^{2} \left[ P_{BFNPZ}^{*} + B_{BFNPZ}^{*} + \alpha + \beta + \delta + k_{24} + k_{26} - \gamma - \varepsilon - 1 \right] + N_{BFNPZ}^{*} \left[ k_{24} k_{26} + \left( k_{24} + k_{26} \right) \left( P_{BFNPZ}^{*} + B_{BFNPZ}^{*} + \alpha - \gamma - \varepsilon - 1 \right) + k_{24} \beta + k_{26} \delta \right]$$

$$+ k_{24} k_{26} \left( P_{BFNPZ}^{*} + B_{BFNPZ}^{*} + \alpha - \gamma - \varepsilon - 1 \right) = 0$$

$$\Rightarrow N_{BFNPZ}^{*} = 0.4707,$$

$$P_{BFNPZ}^* = \frac{k_{19}}{k_4 \left( 1 - k_{20} \right)} = 0.1667, \tag{75}$$

$$Z_{BFNPZ}^* = \left(\frac{k_{23}}{k_4}\right) \left(\frac{N_{BFNPZ}^*}{N_{BFNPZ}^* + k_{24}}\right) - \left(\frac{k_1}{k_4}\right) \left(\frac{B_{BFNPZ}^*}{P_{BFNPZ}^* + k_2}\right) = 0.3394, \quad (76)$$

where 
$$\alpha = k_1 \left( 1 - k_{11} \right) \left( \frac{P_{BFNPZ}^*}{P_{BFNPZ}^* + k_2} \right) \left( \frac{B_{BFNPZ}^* + k_9}{k_8} \right), \qquad \beta = k_{25} \left( 1 - k_{11} \right) \left( \frac{B_{BFNPZ}^* + k_9}{k_8} \right),$$

$$\gamma = k_{10} \left( \frac{B_{BFNPZ}^* + k_9}{k_8} \right)$$
,  $\delta = \frac{k_{23}}{k_4}$  and  $\varepsilon = \frac{k_1}{k_4} \left( \frac{B_{BFNPZ}^*}{P_{BFNPZ}^* + k_2} \right)$  and only the positive root is

considered. The eigenvalues at this point were calculated numerically and are all found to be complex conjugates, with one pair (weakly) unstable.

Figure 1 Click here to download high resolution image




Figure 2




Figure 3 Click here to download high resolution image




Figure 4 Click here to download high resolution image




Figure 5 Click here to download high resolution image




Figure 6 Click here to download high resolution image




Figure 7 Click here to download high resolution image




Figure 8
Click here to download high resolution image




Figure 9 Click here to download high resolution image

